CAMC Prior Authorization Criteria 2020

Effective Date: 12/01/2020 Approval Date: 11/24/2020

#### **Prior Authorization Protocol**

## Medicare Part D - 2020

ACTEMRA SC

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

**Exclusion Criteria:** 

### **Required Medical Information:**

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

RHEUMATOID ARTHRITIS, GIANT CELL ARTERITIS: Prescribed by or in consultation with a rheumatologist. JUVENILE IDIOPATHIC ARTHRITIS: Prescribed by or in consultation with a dermatologist, rheumatologist, or gastroenterologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. GIANT CELL ARTERITIS: Failure of methotrexate or azathioprine, unless contraindicated or clinically significant adverse effects are experienced.

## Medicare Part D – 2020

**Prior Authorization Group Description:** 

day or an equianalgesic dose of another opioid).

ACTIQ

Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
16 years and older.
Prescriber Restrictions:
Coverage Duration:
Through the end of the Plan contract year.
Other Criteria:
Member is already taking and is tolerant to around-the-clock opioid therapy. Members are considered opioid tolerant when taking another opioid daily for a week or longer (for example, at least 60 mg of oral morphine per

# **Medicare Part D – 2020**

<b>Prior Authorization Group Description:</b>
ACYCLOVIR
<b>Prior Authorization Indication:</b>
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
<b>Coverage Duration:</b>
12 months.
Other Criteria:

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**ADAKVEO** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

SICKLE CELL DISEASE: Disease is associated with one of the following genotypes: Homozygous hemoglobin S, Hemoglobin S beta 0-thalassemia, Hemoglobin S beta+ thalassemia, Hemoglobin SC. Member has a hemoglobin level of at least 4 g/dL. Member meets one of the following (a or b): a) Member experienced at least 2 vaso-occlusive crises (VOC) within the past 6 months while on hydroxyurea, OR b) Member has intolerance or contraindication to hydroxyurea and has experienced at least 2 VOC within the past 12 months. Confirmation of baseline incidence of VOC over the last 12 months. Adakveo is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced. Adakveo is not prescribed concurrently with Oxbryta. CONTINUATION OF THERAPY, SICKLE CELL DISEASE: Member is responding positively to therapy as evidenced by an improvement in the incidence of VOC from baseline. Adakveo is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced. Adakveo is not prescribed concurrently with Oxbryta.

#### **Age Restrictions:**

SICKLE CELL DISEASE: Age greater than or equal to 16 years.

#### **Prescriber Restrictions:**

SICKLE CELL DISEASE: Prescribed by or in consultation with a hematologist.

#### **Coverage Duration:**

6 months.

#### Other Criteria:

SICKLE CELL DISEASE: Failure of L-glutamine, unless contraindicated or clinically significant adverse effects are experienced.

# **Medicare Part D – 2020**

<b>Prior Authorization Group Description:</b>
ADCIRCA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Members on concomitant nitrates (e.g., Nitrodur, Nitrobid, Nitrostat, Isordil, Ismo). Members on concomitant guanylate cyclase stimulator, such as riociguat (Adempas).
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

# **Medicare Part D – 2020**

**Prior Authorization Group Description:** 

ADEMPAS
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Members on concomitant phosphodiesterase (PDE) inhibitors (e.g., sildenafil, tadalafil, vardenafil, dipyridamole or theophylline) or nitrates (e.g., Nitrodur, Nitrobid, Nitrostat, Isordil, Ismo).
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**AFINITOR** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

BREAST CANCER: hormone-receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative recurrent or metastatic disease. TUBEROUS SCLEROSIS COMPLEX WITH SUBEPENDYMAL GIANT CELL ASTROCYTOMA: Member is not a candidate for curative surgical resection.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist. TUBEROUS SCLEROSIS COMPLEX ASSOCIATED PARTIAL ONSET SEIZURES: Prescribed by or in consultation with an oncologist or neurologist.

#### **Coverage Duration:**

12 months.

### Other Criteria:

RENAL CELL CARCINOMA WITH CLEAR CELL HISTOLOGY: Failure of one prior therapy (e.g., Votrient, Sutent), unless contraindicated or clinically significant adverse effects are experienced. BREAST CANCER: Prescribed in combination with exemestane, fulvestrant or tamoxifen AND history of prior endocrine therapy (e.g., letrozole, anastrozole) unless contraindicated or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**AIMOVIG** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

One of the following (a or b): a) Member experiences 4 or more migraine days per month for at least 3 months, or b) Diagnosis of chronic migraine and Aimovig is prescribed for prophylaxis. CONTINUATION OF THERAPY: Member has experienced and maintained positive response to therapy as evidenced by a reduction in migraine days per month from baseline.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist, headache, or pain specialist.

#### **Coverage Duration:**

Initial: 3 months. Reauthorizations: 6 months.

### Other Criteria:

Failure of two of the following oral migraine preventative therapies, each from different therapeutic classes, unless contraindicated or clinically significant adverse effects are experienced: antiepileptic drugs (e.g., divalproex sodium, sodium valproate, topiramate), beta-blockers (e.g., metoprolol, propranolol, timolol), antidepressants (e.g., amitriptyline, venlafaxine).

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**AJOVY** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

One of the following (a or b): a) Member experiences 4 or more migraine days per month for at least 3 months, or b) Diagnosis of chronic migraine and Ajovy is prescribed for prophylaxis. CONTINUATION OF THERAPY: Member has experienced and maintained positive response to therapy as evidenced by a reduction in migraine days per month from baseline.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist, headache, or pain specialist.

#### **Coverage Duration:**

Initial: 3 months. Reauthorizations: 6 months.

### Other Criteria:

Failure of two of the following oral migraine preventative therapies, each from different therapeutic classes, unless contraindicated or clinically significant adverse effects are experienced: antiepileptic drugs (e.g., divalproex sodium, sodium valproate, topiramate), beta-blockers (e.g., metoprolol, propranolol, timolol), antidepressants (e.g., amitriptyline, venlafaxine).

## Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>	

**ALECENSA** 

## **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

## Off Label Uses:

## **Exclusion Criteria:**

## **Required Medical Information:**

Disease is anaplastic lymphoma kinase (ALK) positive.

## **Age Restrictions:**

## **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

## **Coverage Duration:**

12 months.

## Medicare Part D – 2020

ALLI
Prior Authorization Indication:
Weight loss in overweight adults, when used along with a reduced-calorie and low-fat diet.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
BMI is greater than or equal to 30 kg/m <sup>2</sup> OR BMI is greater than or equal to 27 kg/m <sup>2</sup> with one or more of the following severe co-morbid conditions 1. Coronary artery/heart disease 2. Diabetes 3. Dyslipidemia 4. Hypertension 5. Obstructive sleep apnea. CONTINUATION OF THERAPY: Confirmation of a 5-10 pound weight loss during the first 6 month treatment period. Therapy beyond the first year can be authorized every 6 months with confirmation of weight maintenance.
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:

6 months.

## Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>		

## **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

## Off Label Uses:

ALUNBRIG

## **Exclusion Criteria:**

## **Required Medical Information:**

Disease is anaplastic lymphoma kinase (ALK) positive.

## **Age Restrictions:**

## **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

## **Coverage Duration:**

12 months.

#### Medicare Part D - 2020

## **Prior Authorization Group Description:**

**AMITRIPTYLINE** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

### **Required Medical Information:**

### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

#### **Prescriber Restrictions:**

#### **Coverage Duration:**

12 months.

## Other Criteria:

Depression: Failure of one of the following generic antidepressants, unless contraindicated or clinically significant adverse effects are experienced: bupropion, bupropion SR, bupropion XL, citalopram, desvenlafaxine succinate, duloxetine, escitalopram, fluoxetine, mirtazapine, sertraline, venlafaxine, or venlafaxine XR.

#### Medicare Part D - 2020

### **Prior Authorization Group Description:**

AMITRIPTYLINE/CHLORDIAZEPOXIDE

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

## **Required Medical Information:**

#### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

#### **Prescriber Restrictions:**

## **Coverage Duration:**

12 months.

## Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: duloxetine, escitalopram, or venlafaxine XR.

## Medicare Part D - 2020

#### **Prior Authorization Group Description:**

AMITRIPTYLINE/PERPHENAZINE

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

## **Required Medical Information:**

#### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

#### **Prescriber Restrictions:**

## **Coverage Duration:**

12 months.

### Other Criteria:

Depression: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: duloxetine, escitalopram, or venlafaxine XR.

## Medicare Part D – 2020

**Prior Authorization Group Description:** 

clinically significant adverse effects are experienced.

AMPHOTERICIN B
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Abelcet only: Failure of conventional amphotericin B therapy unless contraindicated or clinically significant adverse effects are experienced. Ambisome when treating patients with Aspergillus species, Candida species and/or Cryptococcus species infections: Failure of conventional amphotericin B therapy unless contraindicated or

# **Medicare Part D – 2020**

Prior Authorization Group Description:
AMPYRA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a neurologist.
Coverage Duration:
12 months.
Other Criteria:

#### Medicare Part D - 2020

## **Prior Authorization Group Description:**

**ANTIHISTAMINES** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

## **Required Medical Information:**

### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

#### **Prescriber Restrictions:**

### **Coverage Duration:**

12 months.

## Other Criteria:

Allergic rhinitis: Failure to two of the following, unless contraindicated or clinically significant adverse effects are experienced: levocetirizine, desloratadine, fluticasone propionate nasal suspension, flunisolide nasal solution or triamcinolone acetonide nasal inhaler.

#### Medicare Part D - 2020

### **Prior Authorization Group Description:**

ANTIHISTAMINE COMBINATIONS

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

## **Required Medical Information:**

#### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

#### **Prescriber Restrictions:**

### **Coverage Duration:**

12 months.

## Other Criteria:

Allergic rhinitis: Failure to two of the following, unless contraindicated or clinically significant adverse effects are experienced: levocetirizine, desloratadine, fluticasone propionate nasal suspension, flunisolide nasal solution or triamcinolone acetonide nasal inhaler.

# **Medicare Part D – 2020**

Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Anemia due to myelodysplastic syndrome. Myelofibrosis-associated anemia.
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of Procrit, unless contraindicated or clinically significant adverse effects are experienced.

**Prior Authorization Group Description:** 

ARANESP

## Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**ARIKAYCE** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Positive sputum culture after at least 6 consecutive months of a multidrug background regimen therapy (e.g., clarithromycin or azithromycin, ethambutol, and a rifamycin). CONTINUATION OF THERAPY: Confirmation of at least 3 consecutive negative monthly sputum cultures in the first 6 months of therapy or at least 2 consecutive negative monthly sputum cultures in the last 2 months of therapy. Member has not received Arikayce treatment for more than 12 months after converting to negative sputum status.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an infectious disease specialist or pulmonologist.

#### **Coverage Duration:**

Initial: 6 months. Reauthorizations: 12 months.

# Medicare Part D – 2020

Prior Authorization Group Description:
AUBAGIO
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a neurologist.
Coverage Duration:
12 months.
Other Criteria:

## Medicare Part D - 2020

#### **Prior Authorization Group Description:**

AYVAKIT

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

## **Required Medical Information:**

GASTROINTESTINAL STROMAL TUMOR: Confirmation of PDGFRA exon 18 mutation.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

## **Coverage Duration:**

12 months.

### Other Criteria:

GASTROINTESTINAL STROMAL TUMOR: If disease is positive for a PDGFRA exon 18 mutation other than D842V, failure of imatinib, unless contraindicated or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**BALVERSA** 

### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

## **Required Medical Information:**

Presence of susceptible fibroblast growth factor receptor (FGFR) 3 or FGFR2 genetic alterations.

## **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

## **Coverage Duration:**

12 months.

### Other Criteria:

Disease has progressed during or following at least one line of platinum-containing chemotherapy.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**BAXDELA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

ACUTE BACTERIAL SKIN AND SKIN STRUCTURE INFECTION (ABSSSI), COMMUNITY-ACQUIRED BACTERIAL PNEUMONIA (CABP): Request is for continuation of therapy initiated in an acute care hospital from which member was discharged OR Both of the following (1 and 2): 1) Current culture and sensitivity (C&S) report shows isolated pathogen is susceptible to delafloxacin, unless provider confirms that obtaining a C&S report is not feasible AND 2) Failure of one fluoroquinolone, unless all are contraindicated or clinically significant adverse effects are experienced, or C&S report shows resistance or lack of susceptibility of the isolated pathogen to all fluoroquinolones.

### **Age Restrictions:**

#### **Prescriber Restrictions:**

### **Coverage Duration:**

ABSSSI: 14 days. CABP: 10 days.

# Medicare Part D – 2020

<b>Prior Authorization Group Description:</b>
BELEODAQ
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with an oncologist or hematologist.
Coverage Duration:
12 months.
Other Criteria:

## Medicare Part D – 2020

Prior Authorization Group Description:
BELSOMRA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
For patients 65 years of age and older: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Rozerem, Silenor 6 mg/day or less, or trazodone. For patients under 65 years of age: Failure of zolpidem or zolpidem CR, unless contraindicated or clinically significant adverse effects are

experienced.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**BENLYSTA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Confirmation that member is positive for autoantibody (e.g., anti-nuclear antibody (ANA), anti-double-stranded DNA (anti-ds-DNA), anti-Smith antigen (anti-Sm), anti-ribonucleoprotein (anti-RNP), anti-Ro/SSA, anti-La/SSB).

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a rheumatologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

Currently receiving standard therapy for systemic lupus erythematosus that includes one or more of the following agents, unless all agents are contraindicated: glucocorticoids (e.g., prednisone), antimalarial (e.g., hydroxychloroquine or chloroquine), non-biologic immunosuppressants (e.g., azathioprine, methotrexate, mycophenolate).

## Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>		

# Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

## Off Label Uses:

BENZTROPINE

#### **Exclusion Criteria:**

## **Required Medical Information:**

#### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

#### **Prescriber Restrictions:**

## **Coverage Duration:**

12 months.

## Other Criteria:

Parkinsons disease/Parkinsonism: Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: amantadine, levodopa/carbidopa, entacapone, pramipexole, ropinirole, selegiline.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**BEOVU** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

CONTINUATION OF THERAPY, NEOVASCULAR (WET) AGE-RELATED MACULAR DEGENERATION (AMD): Member is responding positively to therapy as evidenced by one of the following (a, b, c, or d): a) detained neovascularization, b) improvement/stabilization in visual acuity, c) maintenance of corrected visual acuity from prior treatment, or d) supportive findings from optical coherence tomography or fluorescein angiography.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

NEOVASCULAR (WET) AMD: Prescribed by or in consultation with an ophthalmologist.

#### **Coverage Duration:**

NEOVASCULAR (WET) AMD: Initial: 4 months. Continuation: 6 months.

### Other Criteria:

NEOVASCULAR (WET) AMD: Failure of intravitreal bevacizumab, unless contraindicated or clinically significant adverse effects are experienced.

# **Medicare Part D – 2020**

Prior Authorization Group Description:
BLEOMYCIN
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with an oncologist.
Coverage Duration:
12 months.
Other Criteria:

## Medicare Part D - 2020

Authorization Group Description:

**BOSULIF** 

## **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

## Off Label Uses:

## **Exclusion Criteria:**

## **Required Medical Information:**

Confirmation that the member has Philadelphia chromosome positive disease.

## **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

## **Coverage Duration:**

12 months.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**BOTOX** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

CHRONIC MIGRAINE HEADACHE: Persistent history of chronic, debilitating migraine headaches with frequent attacks on more than 15 days per month.

#### **Age Restrictions:**

Strabismus or blepharospasm associated with dystonia: 12 years of age or older.

#### **Prescriber Restrictions:**

Chronic migraine headache: Prescribed by or in consultation with a neurologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

Chronic migraine headache: Failure of prophylactic treatment with ONE of the following, unless contraindicated or clinically significant adverse effects are experienced: divalproex, topiramate, timolol or propranolol AND Failure of abortive therapy with ONE of the following, unless contraindicated or clinically significant adverse effects are experienced: sumatriptan, rizatriptan, zolmitriptan, naratriptan, almotriptan, frovatriptan, Relpax, ergotamine/caffeine or dihydroergotamine.

## Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**BRAFTOVI** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

## **Required Medical Information:**

Positive for BRAF V600E or V600K mutation.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

## **Coverage Duration:**

12 months.

## Other Criteria:

MELANOMA: Prescribed in combination with Mektovi. COLON CANCER, RECTAL CANCER: Prescribed in combination with either Erbitux or Vectibix.

## Medicare Part D – 2020

Prior Authorization Group Description:
BRIVIACT
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of two of the following generic antiepileptic drugs, unless contraindicated or clinically significant advers effects are experienced: lamotrigine, topiramate, oxcarbazepine, carbamazepine, phenytoin, valproic acid or

divalproex sodium.

## Medicare Part D - 2020

## **Prior Authorization Group Description:**

**BRUKINSA** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

## Off Label Uses:

#### **Exclusion Criteria:**

## **Required Medical Information:**

#### **Age Restrictions:**

## **Prescriber Restrictions:**

MANTLE CELL LYMPHOMA: Prescribed by or in consultation with an oncologist or hematologist.

## **Coverage Duration:**

12 months.

#### Other Criteria:

MANTLE CELL LYMPHOMA: Previously received at least one prior therapy (e.g., rituximab-containing regimen).

# Medicare Part D - 2020

Prior Authorization	Group	<b>Description:</b>
---------------------	-------	---------------------

C1 ESTERASE INHIBITOR

## **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

# **Required Medical Information:**

Member is not using the requested product in combination with another FDA-approved product for the same indication (e.g., using both Cinryze and Haegarda for long-term prophylaxis of HAE attacks).

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an immunologist, allergist, or hematologist.

#### **Coverage Duration:**

12 months.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**CABLIVI** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Prescribed in combination with plasma exchange therapy. Prescribed in combination with immunosuppressive therapy (i.e., glucocorticoids, rituximab). CONTINUATION OF THERAPY: Member has received no more than 58 days of Cablivi therapy after completion of plasma exchange therapy AND member meets one of the following (a or b): a) If request is for a new treatment cycle, member has experienced no more than two recurrences while taking Cablivi and Cablivi is prescribed in combination with plasma exchange and immunosuppressive therapy (i.e., glucocorticoids, rituximab) OR b) If request is for treatment extension, member is responding positively to therapy as evidenced by, including but not limited to, improvement in any of the following parameters: increase in platelet counts, reduction in neurological symptoms, or improvements in organ-damage markers (lactate dehydrogenase, cardiac troponin I, and serum creatinine).

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a hematologist.

## **Coverage Duration:**

Initial: 60 days. Reauthorization: 58 days post plasma-exchange.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**CABOMETYX** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

# **Required Medical Information:**

NON-SMALL CELL LUNG CANCER: Confirmation of an RET gene rearrangement.

# **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with oncologist.

# **Coverage Duration:**

12 months.

## Other Criteria:

HEPATOCELLULAR CARCINOMA: Failure of Nexavar, unless contraindicated or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

CALQUENCE

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

# **Required Medical Information:**

CLL/SLL: If disease is refractory to Imbruvica, member does not have BTK C481S mutations.

# **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

# **Coverage Duration:**

12 months.

# Other Criteria:

MCL: Previously received at least one prior therapy (e.g., rituximab-containing regimen).

Prior Authorization Group Description:
CAPLYTA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
SCHIZOPHRENIA: Failure of two of the following atypical antipsychotics, unless contraindicated or clinically significant adverse effects are experienced: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole.

# Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**CAPRELSA** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

# **Required Medical Information:**

NON-SMALL CELL LUNG CANCER: Confirmation of an RET gene rearrangement.

# **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# Other Criteria:

DIFFERENTIATED THYROID CARCINOMA: Failure of Lenvima or Nexavar, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:
CAYSTON
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D = 2020

Medicare Part D – 2020
Prior Authorization Group Description:
CERDELGA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Extensive metabolizer (EM) or intermediate metabolizer (IM) taking a strong or moderate CYP2D6 inhibitor concomitantly with a strong or moderate CYP3A inhibitor AND IMs or poor metabolizer (PM) taking a strong CYP3A inhibitor.

# **Required Medical Information:**

An FDA-cleared genotyping test has determined that this patient is a CYP2D6 extensive metabolizer (EM), intermediate metabolizer (IM), or poor metabolizer (PM).

# intermediate metabolizer (IM), or poor metabolizer (PM). Age Restrictions:

# **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

<b>Prior Authorization Group Description:</b>
CEREZYME
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Type 3 Gaucher disease.
Exclusion Criteria:
Required Medical Information:
Confirmation of at least one of the following conditions resulting from Gaucher disease: anemia, thrombocytopenia, bone disease, hepatomegaly or splenomegaly.
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

Prior Authorization Group Description:
CHLORZOXAZONE
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D – 2020

Prior Authorization Group Description:
CHORIONIC GONADOTROPIN
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Induction of ovulation and pregnancy in the anovulatory, infertile woman in whom the cause of anovulation is secondary and not due to primary ovarian failure. Treatment of obesity.
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**CINQAIR** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Blood eosinophil count of greater than or equal to 400 cells/mcL within the past 3 months.

#### **Age Restrictions:**

18 years of age or older.

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an allergist, pulmonologist, or immunologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

Prescribed in combination with ONE inhaled corticosteroid (e.g., beclomethasone, budesonide, flunisolide, fluticasone, mometasone, ciclesonide), unless contraindicated or clinically significant adverse effects are experienced. AND Prescribed in combination with ONE long-acting beta-agonist (e.g., salmeterol, formoterol, vilanterol), unless contraindicated or clinically significant adverse effects are experienced.

<b>Prior Authorization Group Description:</b>
CLADRIBINE
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with an oncologist or hematologist.
<b>Coverage Duration:</b>
12 months.
Other Criteria:

#### Medicare Part D - 2020

Prior Authorization Group Description:

**CLOMIPRAMINE** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

Autistic disorder.

#### **Exclusion Criteria:**

## **Required Medical Information:**

#### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

## **Prescriber Restrictions:**

#### **Coverage Duration:**

12 months.

# Other Criteria:

Failure of one selective serotonin reuptake inhibitor (e.g., fluoxetine, fluvoxamine, sertraline), unless contraindicated or clinically significant adverse effects are experienced.

# Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**COMETRIQ** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

#### **Exclusion Criteria:**

# **Required Medical Information:**

NON-SMALL CELL LUNG CANCER: Confirmation of an RET gene rearrangement.

# **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# Other Criteria:

DIFFERENTIATED THYROID CARCINOMA: Failure of Lenvima or Nexavar unless contraindicated or clinically significant adverse effects are experienced.

<b>Prior Authorization Group Description:</b>
COPIKTRA
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with an oncologist or hematologist.
<b>Coverage Duration:</b>
12 months.
Other Criteria:

# **Medicare Part D – 2020**

# **Prior Authorization Group Description:**

**COTELLIC** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

# Off Label Uses:

#### **Exclusion Criteria:**

Patients with wild-type BRAF melanoma.

#### **Required Medical Information:**

Disease is positive for the BRAF V600E or V600K mutation.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# Other Criteria:

Prescribed in combination with Zelboraf.

# Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>		

## **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

**CRYSVITA** 

#### **Exclusion Criteria:**

## **Required Medical Information:**

X-LINKED HYPOPHOSPHATEMIA: DNA testing results confirm the presence of mutations in the PHEX gene or documentation of elevated serum fibroblast growth factor 23 (FGF23) levels. Current (within the last 30 days) serum phosphorus level is below the reference range for age and gender. CONTINUATION OF THERAPY: Member meets all approval criteria and has had an increase in serum phosphorus level from baseline and/or maintenance within the normal range for age and gender, while on Crysvita therapy.

#### **Age Restrictions:**

At least 6 months of age.

#### **Prescriber Restrictions:**

X-LINKED HYPOPHOSPHATEMIA: Prescribed by or in consultation with an endocrinologist or metabolic disease specialist.

#### **Coverage Duration:**

12 months.

Prior Authorization Group Description:
CYCLOBENZAPRINE HCL
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D – 2020

**Prior Authorization Group Description:** 

CYTARABINE

Prior Authorization Indication:	
All medically accepted indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	
Required Medical Information:	
Age Restrictions:	
Prescriber Restrictions:	
Prescribed by or in consultation with an oncologist or hematologist.	
Coverage Duration:	
2 months.	
Other Criteria:	

For acute non-lymphocytic leukemia: use in combination with other approved anti-cancer drugs.

# Medicare Part D - 2020

Prior Authorization Group Description:	
--	--

**DAURISMO** 

## **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

# **Required Medical Information:**

Age 75 years or greater, OR medical justification supports inability to use intensive induction chemotherapy OR member responded to then relapsed after Daurismo induction therapy 12 or more months ago. Prescribed in combination with low-dose cytarabine.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

## **Coverage Duration:**

12 months.

<b>Prior Authorization Group Description:</b>
DICLOFENAC GEL
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
6 months.
Other Criteria:

#### Medicare Part D - 2020

<b>Prior Authorization</b>	Group	<b>Description:</b>
----------------------------	-------	---------------------

DIETHYLPROPION

# **Prior Authorization Indication:**

The management of exogenous obesity as a short-term adjunct (a few weeks) in a regimen of weight reduction based on caloric restriction in patients with an initial body mass index (BMI) of 30 kg/m<sup>2</sup> or higher and who have not responded to appropriate weight reducing regimen (diet and/or exercise) alone.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

BMI is greater than or equal to 30 kg/m<sup>2</sup> OR BMI is greater than or equal to 27 kg/m<sup>2</sup> with one or more of the following severe co-morbid conditions 1. Coronary artery/heart disease 2. Diabetes 3. Dyslipidemia 4. Hypertension 5. Obstructive sleep apnea.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

## **Coverage Duration:**

Three months.

Prior Authorization Group Description:
DIPYRIDAMOLE
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

Prior Authorization Group Description:
DISOPYRAMIDE
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**DOPTELET** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

THROMBOCYTOPENIA WITH CHRONIC LIVER DISEASE: Recent (within the past 14 days) platelet count is less than 50 x 10^9/L. Member is scheduled to undergo a medical or dental procedure within the next 30 days.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

THROMBOCYTOPENIA WITH CHRONIC LIVER DISEASE: Prescribed by or in consultation with a hematologist, hepatologist, or gastroenterologist. CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): Prescribed by or in consultation with a hematologist.

#### **Coverage Duration:**

THROMBOCYTOPENIA WITH CHRONIC LIVER DISEASE: 4 weeks. CHRONIC ITP: 12 months.

## Other Criteria:

THROMBOCYTOPENIA WITH CHRONIC LIVER DISEASE: For members with platelet count less than 40 x 10^9/L, failure of Mulpleta unless contraindicated or clinically significant adverse effects are experienced. CHRONIC ITP: Failure of a corticosteroid (e.g., prednisone, methylprednisolone, or dexamethasone), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:
DOXEPIN
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

Prior Authorization Group Description:
DOXEPIN CREAM
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
4 weeks.
Other Criteria:
Failure of two topical corticosteroids (e.g., amcinonide, fluticasone propionate, triamcinolone acetonide, betamethasone valerate, fluocinolone acetonide, hydrocortisone butyrate, mometasone furoate, desoximetaso fluocinonide or betamethasone dipropionate), unless contraindicated or clinically significant adverse effects a experienced.

Prior Authorization Group Description:
ELIDEL
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of two medium to high potency topical corticosteroids (e.g., amcinonide, fluticasone propionate, triamcinolone acetonide, betamethasone valerate, fluocinolone acetonide, hydrocortisone butyrate, mometasone furoate, desoximetasone, fluocinonide or betamethasone dipropionate), unless contraindicated or clinically significant adverse effects are experienced.

<b>Prior Authorization Group Description:</b>
EMEND 40 MG
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
4 weeks.
Other Criteria:

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**EMFLAZA** 

## **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

# **Required Medical Information:**

Diagnosis of Duchenne muscular dystrophy (DMD) confirmed by one of the following: Genetic testing (e.g., dystrophin deletion or duplication mutation found) OR if genetic studies are negative (i.e., no mutation identified), positive muscle biopsy (e.g., absence of dystrophin protein).

#### **Age Restrictions:**

2 years of age or older.

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist.

## **Coverage Duration:**

12 months.

## Other Criteria:

Failure of prednisone, unless contraindicated or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**EMGALITY** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

MIGRAINE PROPHYLAXIS: One of the following (a or b): a) Member experiences 4 or more migraine days per month for at least 3 months, or b) Diagnosis of chronic migraine and Emgality is prescribed for prophylaxis. EPISODIC CLUSTER HEADACHE: Member has had at least 2 cluster headache attack periods which lasted for 1 year or less each and were separated by at least 3 months. CONTINUATION OF THERAPY, MIGRAINE PROPHYLAXIS: Member has experienced and maintained positive response to therapy as evidenced by a reduction in migraine days per month from baseline. CONTINUATION OF THERAPY, EPISODIC CLUSTER HEADACHE: Member has experienced and maintained positive response to therapy as evidenced by a reduction in cluster headache attack frequency. Member has not received more than 12 months of consecutive treatment OR it has been at least 3 months since the member last received Emgality.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist, headache, or pain specialist.

## **Coverage Duration:**

Initial: 3 months. Reauthorizations: 6 months.

#### Other Criteria:

MIGRAINE PROPHYLAXIS: Failure of two of the following oral migraine preventative therapies, each from different therapeutic classes, unless contraindicated or clinically significant adverse effects are experienced: antiepileptic drugs (e.g., divalproex sodium, sodium valproate, topiramate), beta-blockers (e.g., metoprolol, propranolol, timolol), antidepressants (e.g., amitriptyline, venlafaxine).

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**ENBREL** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### **Off Label Uses:**

Hidradenitis suppurativa.

#### **Exclusion Criteria:**

#### **Required Medical Information:**

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

PSORIATIC ARTHRITIS, PLAQUE PSORIASIS: Prescribed by or in consultation with a rheumatologist or dermatologist. RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist. HIDRADENITIS SUPPURATIVA: Prescribed by or in consultation with a rheumatologist, dermatologist or gastroenterologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. PLAQUE PSORIASIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin.

Prior Authorization Group Description:
ENDARI
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Age 5 or older.
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced

# **Medicare Part D – 2020**

**Prior Authorization Group Description:** 

ENTYVIO
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a gastroenterologist.
Coverage Duration:
12 months.
Other Criteria:
Failure of Humira or Remicade, unless contraindicated or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**EPCLUSA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

Criteria will be applied consistent with current AASLD-IDSA guidance.

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Refer to the document, "Recommendations for Testing, Managing, and Treating Hepatitis C," by AASLD-IDSA available at http://www.hcvguidelines.org for drug regimen and duration of treatment based on genotype, treatment status, previous drug regimens used, past medical history and comorbidities.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a gastroenterologist, hepatologist or infectious disease physician.

#### **Coverage Duration:**

12 to 24 weeks based on cirrhosis status, genotype, prior treatment, or ribavirin eligibility.

### Medicare Part D - 2020

### **Prior Authorization Group Description:**

**EPIDIOLEX** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

LENNOX-GASTAUT SYNDROME: will be used as adjunctive therapy with other antiepileptic drugs (e.g., Banzel, clobazam, clonazepam, felbamate, lamotrigine, or topiramate).

### **Age Restrictions:**

Age greater than or equal to 2 years.

### **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist.

### **Coverage Duration:**

12 months.

### Other Criteria:

LENNOX-GASTAUT SYNDROME: Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: Banzel, clobazam, clonazepam, felbamate, lamotrigine, topiramate.

# Medicare Part D – 2020

Prior Authorization Group Description:
EPOETIN
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Anemia due to myelodyspastic syndrome. Anemia associated with myelofibrosis. Anemia secondary to combination ribavirin and interferon-alfa therapy in patients infected with hepatitis C virus.
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D - 2020

Prior Authorization	Group	<b>Description:</b>
---------------------	-------	---------------------

**ERGOLOID MESYLATES** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

### **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

# Other Criteria:

Alzheimer's dementia: Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: donepezil, memantine, rivastigmine or galantamine.

# Medicare Part D - 2020

Prior Authorization Group Description:
--

**ERLEADA** 

### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

PROSTATE CANCER: Concurrent use of a gonadotropin-releasing hormone (GnRH) analog or past bilateral orchiectomy. Disease is characterized as one of the following (a or b): a) not metastatic and castration-resistant, OR b) metastatic and castration-sensitive.

### **Age Restrictions:**

### **Prescriber Restrictions:**

PROSTATE CANCER: Prescribed by or in consultation with an oncologist or urologist.

### **Coverage Duration:**

12 months.

# **Medicare Part D – 2020**

<b>Prior Authorization Group Description:</b>		
ESBRIET		
<b>Prior Authorization Indication:</b>		
All FDA-approved indications not otherwise excluded from Part D.		
Off Label Uses:		
Exclusion Criteria:		
Required Medical Information:		
Age Restrictions:		
Prescriber Restrictions:		
<b>Coverage Duration:</b>		
12 months.		
Other Criteria:		

### Medicare Part D - 2020

### **Prior Authorization Group Description:**

ESTROGENS(Divigel , Climara , Mimvey , Prempro , Amabelz , Estrace , Activella , Femhrt , Mimvey Lo , Lopreeza , Premarin , Premphase )

### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

### **Required Medical Information:**

### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

### **Prescriber Restrictions:**

### **Coverage Duration:**

12 months.

### Other Criteria:

Atrophic Vaginitis and Kraurosis Vulvae: Failure to one of the following, unless contraindicated or clinically significant adverse effects are experienced: Estradiol vaginal tablet, Femring or Premarin vaginal cream.

### Medicare Part D - 2020

### **Prior Authorization Group Description:**

**EXONDYS 51** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

Duchenne muscular dystrophy with mutation amenable to exon 51 skipping confirmed by genetic testing.

### **Age Restrictions:**

### **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist.

# **Coverage Duration:**

6 months.

# Other Criteria:

Currently stable on an oral corticosteroid regimen (e.g., prednisone), unless contraindicated or clinically significant adverse effects are experienced.

# Medicare Part D - 2020

Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:

# **Prescriber Restrictions:**

**Prior Authorization Group Description:** 

Prescribed by or in consultation with a hematologist or oncologist.

# **Coverage Duration:**

12 months.

**FARYDAK** 

### Other Criteria:

Failure of two prior regimens, including bortezomib and an immunomodulatory agent (e.g., dexamethasone), unless contraindicated or clinically significant adverse effects are experienced.

### Medicare Part D - 2020

### **Prior Authorization Group Description:**

**FASENRA** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

### **Required Medical Information:**

Blood eosinophil count of greater than or equal to 150 cells/mcL within the past 3 months.

### **Age Restrictions:**

12 years of age or older.

### **Prescriber Restrictions:**

Prescribed by or in consultation with an allergist, pulmonologist, or immunologist.

### **Coverage Duration:**

12 months.

### Other Criteria:

Prescribed in combination with ONE inhaled corticosteroid (e.g., beclomethasone, budesonide, flunisolide, fluticasone, mometasone, ciclesonide), unless contraindicated or clinically significant adverse effects are experienced AND Prescribed in combination with ONE long-acting beta-agonist (e.g., salmeterol, formoterol, vilanterol), unless contraindicated or clinically significant adverse effects are experienced.

# **Medicare Part D – 2020**

Prior Authorization Group Description:
FERRIPROX
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of deferoxamine. Exiade or Jadenu, unless contraindicated or clinically significant adverse effects are

experienced.

# Medicare Part D – 2020

<b>Prior Authorization Group Description:</b>
FINTEPLA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
DRAVET SYNDROME: Prescribed by or in consultation with a neurologist.
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D - 2020

<b>Prior Authorization Group</b>	Description:		

FIORINAL WITH CODEINE

# **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

### **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

# Other Criteria:

Failure of naproxen and ibuprofen, unless contraindicated or clinically significant adverse effects are experienced.

# Medicare Part D - 2020

**FIRAZYR** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

Member is not using Firazyr in combination with another FDA-approved product for treatment of acute HAE attacks (e.g., Berinert, Ruconest, Kalbitor).

### **Age Restrictions:**

Age 18 or greater.

### **Prescriber Restrictions:**

Prescribed by or in consultation with an immunologist, allergist, or hematologist.

### **Coverage Duration:**

12 months.

# Medicare Part D – 2020

**Prior Authorization Group Description:** 

F.	IRDAPSE
<u>P</u>	rior Authorization Indication:
A	ll FDA-approved indications not otherwise excluded from Part D.
0	off Label Uses:
<u>E</u>	xclusion Criteria:
R	equired Medical Information:
Q (7	onfirmation of a baseline clinical muscle strength assessment (examples may include but are not limited to the puantitative Myasthenia Gravis (QMG) score, triple-timed up-and-go test (3TUG), Timed 25-foot Walk test (25FW)). CONTINUATION OF THERAPY: Member is responding positively to therapy as evidenced by clinical nuscle strength assessments.
<u>A</u>	ge Restrictions:
<u>P</u>	rescriber Restrictions:
P	rescribed by or in consultation with a neurologist.
<u>C</u>	overage Duration:
12	2 months.

# **Medicare Part D – 2020**

**Prior Authorization Group Description:** 

FLECTOR
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Cancer pain.
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
Acute Pain: 4 weeks. Cancer pain: Through the end of the Plan contract year.
Other Criteria:

# Medicare Part D – 2020

Prior Authorization Group Description:	
FLUOROURACIL	
Prior Authorization Indication:	
All medically accepted indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	
Required Medical Information:	
Age Restrictions:	
Prescriber Restrictions:	
Prescribed by or in consultation with an oncologist.	
<b>Coverage Duration:</b>	
12 months.	
Other Criteria:	

### Medicare Part D - 2020

Prior Authorization	<b>Group Description:</b>	

**FORTEO** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

### **Exclusion Criteria:**

### **Required Medical Information:**

Total duration of therapy on parathyroid hormone (PTH) analogs (e.g., Tymlos, Forteo) has not exceeded 2 years.

### **Age Restrictions:**

### **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

#### Other Criteria:

Member meets one of the following (a, b, or c): a) Failure of a bisphosphonate (e.g., alendronate) unless contraindicated or clinically significant adverse effects are experienced. OR b) Bone mineral density T-score at hip or spine is -3.5 or less. OR c) Bone mineral density T-score at hip or spine is -2.5 or less with a history of major osteoporotic fracture of the hip, spine, forearm, wrist, or humerus.

# Medicare Part D - 2020

### **Prior Authorization Group Description:**

**GALAFOLD** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

### **Exclusion Criteria:**

Amenable GLA variants (mutations) associated with benign phenotypes (i.e., phenotypes known not to cause Fabry disease), including the following GLA mutation: c.937G to T, (p.(D313Y)).

### **Required Medical Information:**

Presence of at least one amenable GLA variant (mutation) as confirmed by one of the following resources: Galafold Prescribing Information brochure (package insert - Section 12, Table 2), Amicus Fabry GLA Gene Variant Search Tool: http://www.fabrygenevariantsearch.com/hcp, or Amicus Medical Information at 1-877-4AMICUS or medinfousa@amicusrx.com.

### **Age Restrictions:**

### **Prescriber Restrictions:**

Prescribed by or in consultation with a clinical geneticist.

### **Coverage Duration:**

12 months.

# **Medicare Part D – 2020**

<b>Prior Authorization Group Description:</b>
GANCICLOVIR
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D - 2020

Prior Authorization	<b>Group Description:</b>	

**GATTEX** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

Member has been dependent on parenteral nutrition or other intravenous support for at least 12 months. CONTINUATION OF THERAPY: Requirement for parenteral nutrition or other intravenous support has decreased since initiation of Gattex therapy.

### **Age Restrictions:**

### **Prescriber Restrictions:**

Prescribed by or in consultation with a gastroenterologist.

### **Coverage Duration:**

12 months.

# **Medicare Part D – 2020**

# **Prior Authorization Group Description:**

GAVRETO

**Pending CMS Review** 

# Medicare Part D - 2020

Prior Authoriz	ation Groun	Description:	

**GILENYA** 

# **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

Baseline QTc interval greater than or equal to 500 msec.

# **Required Medical Information:**

# **Age Restrictions:**

10 years of age or older.

# **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist.

# **Coverage Duration:**

12 months.

### Medicare Part D - 2020

**GILOTRIF** 

### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

NON-SMALL CELL LUNG CANCER: Member meets one of the following (a or b): a) disease is positive for a sensitizing EGFR mutation (e.g., exon 19 deletion or insertion, exon 21 point mutation (L858R, L861Q), exon 18 point mutation (G719X), exon 20 point mutation (S768I)), OR b) disease is squamous cell carcinoma histology with progression after platinum-based chemotherapy (e.g., cisplatin, carboplatin).

### **Age Restrictions:**

### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

### **Coverage Duration:**

12 months.

# Medicare Part D – 2020

**Prior Authorization Group Description:** 

GLATIRAMER
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a neurologist.
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>		
GLIMEPIRIDE		

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

### **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

### Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: glipizide or glipizide/metformin combination product.

### Medicare Part D - 2020

# **Prior Authorization Group Description:**

**GLYBURIDE** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

### **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

# Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: glipizide or glipizide/metformin combination product.

### Medicare Part D - 2020

	<b>Prior</b>	Authorization	Group	Descrip	tion:
--	--------------	---------------	-------	---------	-------

GLYBURIDE/METFORMIN

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

### **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

### Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: glipizide or glipizide/metformin combination product.

# Medicare Part D – 2020

GRANIX
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Myelodysplastic syndrome.
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.

**Prior Authorization Group Description:** 

### Medicare Part D - 2020

### **Prior Authorization Group Description:**

**HARVONI** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

### Off Label Uses:

Criteria will be applied consistent with current AASLD-IDSA guidance.

### **Exclusion Criteria:**

### **Required Medical Information:**

Refer to the document, "Recommendations for Testing, Managing, and Treating Hepatitis C," by AASLD-IDSA available at http://www.hcvguidelines.org for drug regimen and duration of treatment based on genotype, treatment status, previous drug regimens used, past medical history and comorbidities.

### **Age Restrictions:**

### **Prescriber Restrictions:**

Prescribed by or in consultation with a gastroenterologist, hepatologist or infectious disease physician.

### **Coverage Duration:**

8 to 24 weeks based on cirrhosis status, genotype, prior treatment, or ribavirin eligibility.

### Medicare Part D - 2020

### **Prior Authorization Group Description:**

**HERCEPTIN** 

### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

### Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

Disease is human epidermal growth factor receptor (HER2) positive.

### **Age Restrictions:**

### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# Other Criteria:

GASTRIC AND ESOPHAGEAL CANCER: prescribed in combination with systemic chemotherapy (e.g., cisplatin and either capecitabine or 5-fluorouracil). ENDOMETRIAL CARCINOMA: prescribed in combination with carboplatin and paclitaxel.

# Medicare Part D – 2020

<b>Prior Authorization Group Description:</b>
HETLIOZ
<b>Prior Authorization Indication:</b>
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
<b>Coverage Duration:</b>
12 months.
Other Criteria:

### Medicare Part D - 2020

### **Prior Authorization Group Description:**

**HUMAN GROWTH HORMONE** 

### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

### **Exclusion Criteria:**

### **Required Medical Information:**

CHILDREN AND ADOLESCENTS WITH GROWTH HORMONE DEFICIENCY, SHOX DEFICIENCY IN CHILDREN: Baseline height must be greater than 2 standard deviations below the mean for gender and age. Growth rate is such that the member is unlikely to attain an adult height in the normal range - 59 inches for girls and 63 inches for boys. TURNER SYNDROME: Confirmed by karyotype. PRADER-WILLI or NOONAN SYNDROME: Baseline height must be less than the 5th percentile for gender and age OR 2 or more standard deviations below the mean measured paternal height. Growth rate is such that the member is unlikely to attain an adult height in the normal range - 59 inches for girls and 63 inches for boys.

### **Age Restrictions:**

### **Prescriber Restrictions:**

### **Coverage Duration:**

Adult Growth Hormone Deficiency: 12 months. HIV Wasting or Cachexia, Children: 6 months.

#### Other Criteria:

HIV Wasting or Cachexia: Member is being treated with concomitant antiretroviral therapy.

### Medicare Part D - 2020

Prior	Authorization	Group	Description:	

**HUMIRA** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

**Exclusion Criteria:** 

### **Required Medical Information:**

**Age Restrictions:** 

### **Prescriber Restrictions:**

For the following indications, prescribed by or in consultation with: PSORIATIC ARTHRITIS, PLAQUE PSORIASIS - rheumatologist or dermatologist. CROHN'S DISEASE, ULCERATIVE COLITIS - gastroenterologist. RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS - rheumatologist. HIDRADENITIS SUPPURATIVA - rheumatologist, dermatologist or gastroenterologist. UVEITIS - ophthalmologist or rheumatologist.

### **Coverage Duration:**

12 months.

### Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. PLAQUE PSORIASIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin.

# Medicare Part D – 2020

Prior Authorization Group Description:	
HYDROCODONE	
Prior Authorization Indication:	
All FDA-approved indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	
Required Medical Information:	
Age Restrictions:	
Prescriber Restrictions:	
Coverage Duration:	
3 months initial for non-malignant pain then 12 months. 12 months for cancer pain.	
Other Criteria:	
Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced:	MS

Contin, Kadian, Duragesic, Opana ER, Avinza or Oxycontin.

# Medicare Part D – 2020

Prior Authorization Group Description:
HYDROXYZINE HCL INJECTION
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

### Medicare Part D - 2020

### **Prior Authorization Group Description:**

HYDROXYZINE HCL ORAL

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

### **Exclusion Criteria:**

### **Required Medical Information:**

### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

### **Prescriber Restrictions:**

### **Coverage Duration:**

12 months.

### Other Criteria:

Pruritus: Failure of one of the following topical agents, unless contraindicated or clinically significant adverse effects are experienced: betamethasone, hydrocortisone, triamcinolone, fluticasone, clobetasol, fluocinonide or fluocinolone. Anxiety: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: venlafaxine, buspirone, duloxetine or escitalopram. All other FDA approved indications: Patient is continuing on this medication without adverse effects.

# Medicare Part D - 2020

# **Prior Authorization Group Description:**

HYDROXYZINE PAMOATE

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

# **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

# **Prescriber Restrictions:**

#### **Coverage Duration:**

12 months.

# Other Criteria:

Pruritus: Failure of one of the following topical agents, unless contraindicated or clinically significant adverse effects are experienced: betamethasone, hydrocortisone, triamcinolone, fluticasone, clobetasol, fluocinonide or fluocinolone. Anxiety: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: venlafaxine, buspirone, duloxetine or escitalopram. All other FDA approved indications: Patient is continuing on this medication without adverse effects.

# Medicare Part D – 2020

ICLUSIG
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
ACUTE LYMPHOBLASTIC LEUKEMIA (ALL): disease is Philadelphia chromosome positive (Ph+).
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with an oncologist or hematologist.

**Prior Authorization Group Description:** 

**Coverage Duration:** 

12 months.

# Medicare Part D - 2020

# **Prior Authorization Group Description:**

**IDHIFA** 

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

Presence of an isocitrate dehydrogenase-2 (IDH2) mutation.

# **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

# **Coverage Duration:**

12 months.

# Other Criteria:

For age less than 60 years, disease has relapsed or is refractory following treatment with a first line chemotherapy regimen (e.g., cytarabine, idarubicin, daunorubicin, Vyxeos, cladribine, Rydapt, Mylotarg).

# Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>		

# **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

# Off Label Uses:

**ILARIS** 

Acute gouty arthritis.

# **Exclusion Criteria:**

# **Required Medical Information:**

Confirmation of current weight.

#### **Age Restrictions:**

Cryopyrin-Associated Periodic Syndromes: 4 years and older. All other covered indications: 2 years and older.

# **Prescriber Restrictions:**

SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS: Prescribed by or in consultation with a dermatologist, rheumatologist, or gastroenterologist. ALL OTHER COVERED INDICATIONS: Prescribed by or in consultation with a rheumatologist.

# **Coverage Duration:**

12 months.

# Medicare Part D - 2020

**Prior Authorization Group Description:** 

All FDA-approved indications not otherwise excluded from Part D.

**Prior Authorization Indication:** 

**ILUMYA** 

Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a rheumatologist or dermatologist.
Coverage Duration:
12 months.
Other Criteria:

Failure of one of the following unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin. Failure of one of the following unless contraindicated or clinically significant adverse effects are experienced: Cosentyx, Humira, Inflectra, Remicade, Stelara, Taltz, and Tremfya.

#### Medicare Part D - 2020

**IMATINIB** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

# **Exclusion Criteria:**

#### **Required Medical Information:**

CHRONIC MYELOID LEUKEMIA (CML), ACUTE LYMPHOBLASTIC LEUKEMIA (ALL): disease is Philadelphia chromosome positive. CHRONIC MYELOMONOCYTIC LEUKEMIA: disease is positive for a platelet-derived growth factor receptor (PDGFR) mutation or a 5q3133 mutation. MYELODYSPLASTIC/MYELOPROLIFERATIVE DISEASES: disease is positive for a PDGFR mutation. AGGRESSIVE SYSTEMIC MASTOCYTOSIS: disease is D816V c-Kit mutation negative or c-Kit mutational status is unknown. MELANOMA: disease is KIT-positive. PIGMENTED VILLONODULAR SYNOVITIS/TENOSYNOVIAL GIANT CELL TUMOR: Disease is associated with severe morbidity or functional limitations and is not amenable to improvement with surgery.

#### **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

### **Coverage Duration:**

12 months.

Prior Authorization Group Description:
IMBRUVICA
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
CHRONIC GRAFT-VERSUS-HOST DISEASE: Prescribed by or in consultation with an oncologist, hematologist or bone marrow transplant specialist. ALL OTHER INDICATIONS: Prescribed by or in consultation with an oncologist or hematologist.
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D - 2020

Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.
Prescriber Restrictions:

# **Coverage Duration:**

**Prior Authorization Group Description:** 

**IMIPRAMINE** 

12 months.

# Other Criteria:

Depression: Failure of one of the following generic antidepressants, unless contraindicated or clinically significant adverse effects are experienced: bupropion, bupropion SR, bupropion XL, citalopram, desvenlafaxine succinate, duloxetine, escitalopram, fluoxetine, mirtazapine, sertraline, venlafaxine or venlafaxine XR.

# Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>	

INDOMETHACIN

# **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

# **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

# **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

# Other Criteria:

Failure of naproxen and sulindac, unless contraindicated or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

# Prior Authorization Group Description:

**INFLECTRA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

**Exclusion Criteria:** 

# **Required Medical Information:**

**Age Restrictions:** 

#### **Prescriber Restrictions:**

Psoriatic Arthritis/Plaque Psoriasis: Prescribed by or in consultation with a rheumatologist or dermatologist. Crohn's Disease/Ulcerative Colitis: Prescribed by or in consultation with a gastroenterologist. Rheumatoid Arthritis/Ankylosing Spondylitis: Prescribed by or in consultation with a rheumatologist.

#### **Coverage Duration:**

12 months.

# Other Criteria:

Rheumatoid Arthritis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. Plaque Psoriasis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin.

# Medicare Part D - 2020

Prior Authorization Group Description:

**INGREZZA** 

# **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

Development of tardive dyskinesia is secondary to a centrally acting dopamine receptor blocking agent (neuroleptic) (e.g., first- or second-generation antipsychotics such as chlorpromazine or aripiprazole, antiemetics such as promethazine or metoclopramide, the tri-cyclic antidepressant amoxapine).

#### **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with a psychiatrist or neurologist.

# **Coverage Duration:**

12 months.

#### Medicare Part D - 2020

# **Prior Authorization Group Description:**

**INLYTA** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

**Exclusion Criteria:** 

#### **Required Medical Information:**

**Age Restrictions:** 

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

#### Other Criteria:

RENAL CELL CARCINOMA WITH CLEAR CELL HISTOLOGY: One of the following (a or b): a) prescribed concurrently with Keytruda or Bavencio, OR b) prescribed as monotherapy after failure of one prior therapy (e.g., Votrient, Sutent), unless contraindicated or clinically significant adverse effects are experienced. DIFFERENTIATED THYROID CARCINOMA: Failure of Lenvima or Nexavar unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:
INQOVI
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist or hematologist.
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D - 2020

or Authorization G	Group Description:
REBIC	

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

MYELOFIBROSIS: Confirmation of a recent (within the last 30 days) thiamine level of 70 nmol/L (3 mcg/dL) or greater. Confirmation of a recent (within the last 30 days) platelet count of 50,000/mcL or greater.

# **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

# **Coverage Duration:**

12 months.

Prior Authorization Group Description:	
INTERFERON BETA-1A	
Prior Authorization Indication:	
All FDA-approved indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	
Required Medical Information:	
Age Restrictions:	
Prescriber Restrictions:	
Prescribed by or in consultation with a neurologist.	
<b>Coverage Duration:</b>	
12 months.	
Other Criteria:	

<b>Prior Authorization Group Description:</b>	
INTERFERON BETA-1B	
Prior Authorization Indication:	
All FDA-approved indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	
Required Medical Information:	
Age Restrictions:	
Prescriber Restrictions:	
Prescribed by or in consultation with a neurologist.	
<b>Coverage Duration:</b>	
12 months.	
Other Criteria:	

# Medicare Part D - 2020

Prior Au	ıthorization	Group	Descri	ption:	

**INTUNIV** 

# **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

# **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

# **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

# Other Criteria:

Attention Deficit Hyperactivity Disorder: Failure of both of the following, unless contraindicated or clinically significant adverse effects are experienced: amphetamine-based stimulant and methylphenidate based-stimulant.

#### Medicare Part D - 2020

# **Prior Authorization Group Description:**

**JAKAFI** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

# **Exclusion Criteria:**

#### **Required Medical Information:**

STEROID REFRACTORY GRAFT-VERSUS-HOST DISEASE: Member has history of bone marrow or stem cell transplant.

#### **Age Restrictions:**

STEROID REFRACTORY GRAFT-VERSUS-HOST DISEASE: Age greater than or equal to 12 years.

# **Prescriber Restrictions:**

STEROID REFRACTORY GRAFT-VERSUS-HOST DISEASE: Prescribed by or in consultation with an oncologist, hematologist, or bone marrow transplant specialist. ALL OTHER INDICATIONS: Prescribed by or in consultation with an oncologist or hematologist.

# **Coverage Duration:**

12 months.

#### Other Criteria:

POLYCYTHEMIA VERA: Failure of hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced. STEROID REFRACTORY GRAFT-VERSUS-HOST DISEASE: Failure of systemic corticosteroids, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:
JUXTAPID
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of Repatha 420 mg, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:
JYNARQUE
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a nephrologist.
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D - 2020

Prior Authorization Group Description:	
KADCYLA	

Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

**Prior Authorization Indication:** 

Kadcyla will be used as a single-agent therapy.

All medically accepted indications not otherwise excluded from Part D.

# **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# **Medicare Part D – 2020**

**Prior Authorization Group Description:** 

KALYDECO
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Presence of one mutation in the CFTR gene that is responsive to ivacaftor. Confirmation that a homozygous F508del mutation in the CFTR gene is not present.
Age Restrictions:
6 months of age or older.
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

KETOROLAC TROMETHAMINE

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

# **Exclusion Criteria:**

Patients with active peptic ulcer disease. Advanced renal impairment or at risk for renal failure due to volume depletion. Suspected or confirmed cerebrovascular bleeding, hemorrhagic diathesis, incomplete hemostasis and those at high risk for bleeding. Patient currently receiving aspirin or NSAIDs (non-steroidal anti-inflammatory drugs). Patient currently receiving probenecid or pentoxifylline.

# **Required Medical Information:**

# **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

# **Prescriber Restrictions:**

# **Coverage Duration:**

5 days.

# Medicare Part D – 2020

**Prior Authorization Group Description:** 

**KEVZARA** 

Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a rheumatologist.
Coverage Duration:
12 months.
Other Criteria:
Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced:

methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin.

#### Medicare Part D - 2020

# **Prior Authorization Group Description:**

KISQALI(Kisqali Femara Co-Pack, Kisqali)

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

# **Exclusion Criteria:**

#### **Required Medical Information:**

Breast cancer is hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, and advanced or metastatic.

#### **Age Restrictions:**

### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

#### **Coverage Duration:**

12 months.

# Other Criteria:

For Kisqali: Prescribed in combination with an aromatase inhibitor (e.g., letrozole, anastrozole, or exemestane), fulvestrant, or tamoxifen. If prescribed in combination with tamoxifen: Medical justification supports need to use tamoxifen over an aromatase inhibitor or fulvestrant. For men receiving an aromatase inhibitor: Prescribed in combination with an agent that suppresses testicular steroidogenesis (e.g., gonadotropin-releasing hormone agonists).

Prior Authorization Group Description:	
KORLYM	
Prior Authorization Indication:	
All FDA-approved indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	
Required Medical Information:	
Age Restrictions:	
Prescriber Restrictions:	
Prescribed by or in consultation with an endocrinologist.	
<b>Coverage Duration:</b>	
12 months.	
Other Criteria:	

# Medicare Part D - 2020

Prior	Authorization	Group	<b>Description:</b>

**KOSELUGO** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

NEUROFIBROMATOSIS TYPE 1 (NF1): Diagnosis is confirmed by positive genetic testing for NF1 or member has at least one diagnostic criteria for NF1 based on the National Institutes of Health Neurofibromatosis 1 Diagnostic Criterion. Complete resection of plexiform neurofibroma (PN) is not considered to be feasible without substantial risk or morbidity (e.g., due to encasement of, or close proximity to, vital structures, invasiveness, or high vascularity of the PN).

# **Age Restrictions:**

#### **Prescriber Restrictions:**

NF1: Prescribed by or in consultation with an oncologist or neurologist.

# **Coverage Duration:**

12 months.

# Medicare Part D - 2020

# **Prior Authorization Group Description:**

**KUVAN** 

# **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

Recent (within 90 days) phenylalanine (Phe) blood level is greater than 360 micromol/L. CONTINUATION OF THERAPY: Confirmation of a reduction in Phe blood levels since initiation of therapy.

#### **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with a metabolic or genetic disease specialist.

# **Coverage Duration:**

Initial: 3 months. Reauthorization: 12 months.

Prior Authorization Group Description:
LATUDA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of two of the following atypical antipsychotics, unless contraindicated or clinically significant adverse effects are experienced: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole.

Prior Authorization Group Description:
LAZANDA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Age 18 or greater
Prescriber Restrictions:
Coverage Duration:
Through the end of the Plan contract year.
Other Criteria:
Patient is already taking and is tolerant to around-the-clock opioid therapy. Patients are considered opioid tolerant when taking another opioid daily for a week or longer (for example, at least 60 mg of oral morphine per day or an equianalgesic dose of another opioid).

Prior Authorization Group Description:
LEMTRADA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a neurologist.
Coverage Duration:
12 months.
Other Criteria:
Failure of TWO of the following, unless contraindicated or clinically significant adverse effects are experienced Aubagio, Tecfidera, Gilenya, Avonex, Betaseron, Plegridy, glatiramer, Copaxone, Glatopa, Extavia or Rebif.

#### Medicare Part D - 2020

Prior Authorization Group Description	1.		

LENVIMA

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

**Exclusion Criteria:** 

#### **Required Medical Information:**

**Age Restrictions:** 

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

#### Other Criteria:

RENAL CELL CARCINOMA: Prescribed in combination with Afinitor AND if histology is clear cell or unknown, failure of a regimen consisting of or including one of the following drugs unless contraindicated or clinically significant adverse effects are experienced: Avastin, Cabometyx, Inlyta, Nexavar, Opdivo, Proleukin, Sutent, Tarceva, Torisel, Votrient, or Yervoy. MEDULLARY THYROID CARCINOMA: Failure of Cometriq or Caprelsa unless contraindicated or clinically significant adverse effects are experienced.

# Medicare Part D - 2020

Prior Authorization Group Description:
LEUKINE
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:

# **Other Criteria:**

12 months.

AML following induction therapy, Mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation, Following autologous peripheral blood progenitor cell or bone marrow transplantation (BMT) in members with NHL, ALL, HL for acceleration of myeloid reconstitution, Following allogeneic BMT for acceleration of myeloid reconstitution, Acute Radiation Syndrome: Failure of Neupogen, unless contraindicated or clinically significant adverse effects are experienced.

# **Medicare Part D - 2020**

**Prior Authorization Group Description:** 

All FDA-approved indications not otherwise excluded from Part D.

Diabetic neuropathy. Cancer-related neuropathic pain.

**Prior Authorization Indication:** 

LIDODERM

Off Label Uses:

**Exclusion Criteria:** 

Required Medical Information:		
Age Restrictions:		
Prescriber Restrictions:		
Coverage Duration:		
12 months.		
Other Criteria:		

#### Medicare Part D - 2020

# **Prior Authorization Group Description:**

LONSURF

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

# **Exclusion Criteria:**

#### **Required Medical Information:**

COLORECTAL CANCER: Confirmation that the patient does or does not have the RAS (KRAS or NRAS) wild type gene. GASTRIC CANCER, GASTROESOPHAGEAL ADENOCARCINOMA: Confirmation that the patient does or does not have a HER2/neu-positive tumor.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

### **Coverage Duration:**

12 months.

#### Other Criteria:

COLORECTAL CANCER: If tumor expresses the RAS wild type gene, failure of Erbitux or Vectibix, unless contraindicated or clinically significant adverse effects are experienced. GASTRIC CANCER, GASTROESOPHAGEAL ADENOCARCINOMA: If tumor is HER2/neu-positive (i.e., HER2-overexpressing), failure of Herceptin, unless contraindicated or clinically significant adverse effects are experienced.

# Medicare Part D - 2020

# **Prior Authorization Group Description:**

**LORBRENA** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

NON-SMALL CELL LUNG CANCER: Disease is ALK or ROS1 positive.

# **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# Other Criteria:

NON-SMALL CELL LUNG CANCER: For ALK-positive disease, failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Alecensa, Alunbrig, Xalkori, Zykadia. For ROS1-positive disease, failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Rozlytrek, Xalkori, Zykadia.

### Medicare Part D – 2020

**Prior Authorization Group Description:** 

All FDA-approved indications not otherwise excluded from Part D.

**Prior Authorization Indication:** 

LOTRONEX

Off Label Uses:

**Exclusion Criteria:** 

Male patients.
Required Medical Information:
Female patient with irritable bowel symptoms persisting for at least 6 months.
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**LUCEMYRA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Diagnosis of opioid dependence (may be limited to physiologic dependence/tolerance) or opioid use disorder. Member is currently or will be undergoing abrupt opioid discontinuation within the next seven days and one of the following: member has taken one or more opioids for at least the last three weeks OR an opioid antagonist (e.g., naltrexone) has been or will be administered after a period of opioid use. Medical justification supports why an opioid taper (e.g., with buprenorphine, methadone or other opioid) cannot be used. Lucemyra has not been prescribed for a prior opioid withdrawal event within the last 30 days or medical justification supports retreatment.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a physician specializing in one of the following areas: emergency medicine/inpatient care, pain management, addiction psychiatry.

#### **Coverage Duration:**

14 days per course of treatment.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

LYNPARZA TABLET

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

BREAST CANCER: Mutations in the BRCA genes and confirmation of human epidermal growth factor receptor 2 (HER2)-negative disease. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: Member does not have a PPP2R2A gene mutation.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

PROSTATE CANCER: Prescribed by or in consultation with an oncologist or urologist. ALL OTHER ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

#### **Coverage Duration:**

12 months.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**MAVENCLAD** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

CONTINUATION OF THERAPY: Member has not yet received 2 courses (4 cycles) lifetime total.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

RELAPSING-REMITTING MULTIPLE SCLEROSIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Aubagio, Tecfidera, Gilenya, Avonex, Betaseron, Plegridy, glatiramer, Copaxone, Glatopa, Extavia, or Rebif.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**MAVYRET** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

Treatment-experienced patients with both NS3/4A protease inhibitor and NS5A inhibitor.

#### **Required Medical Information:**

If cirrhosis is present, confirmation of Child-Pugh A status. Refer to the document, "Recommendations for Testing, Managing, and Treating Hepatitis C," by AASLD-IDSA available at http://www.hcvguidelines.org for drug regimen and duration of treatment based on genotype, treatment status, previous drug regimens used, past medical history and comorbidities.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a gastroenterologist, hepatologist or infectious disease physician.

#### **Coverage Duration:**

8 to 16 weeks based on genotype, cirrhosis status, prior treatment regimen.

#### Other Criteria:

If member has been previously treated with an HCV regimen containing NS5A inhibitor or an NS3/4A protease inhibitor, but not both, member has genotype 1.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**MAYZENT** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

CYP2C9\*3/\*3 genotype.

#### **Required Medical Information:**

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

RELAPSING-REMITTING MULTIPLE SCLEROSIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Aubagio, Tecfidera, Gilenya, Avonex, Betaseron, Plegridy, glatiramer, Copaxone, Glatopa, Extavia, or Rebif.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**MEGACE** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

BREAST CANCER AND ENDOMETRIAL CANCER: Megestrol acetate is being used for palliative treatment.

#### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

#### **Prescriber Restrictions:**

#### **Coverage Duration:**

12 months.

#### Other Criteria:

ANOREXIA AND CACHEXIA ASSOCIATED WITH AIDS: Failure of dronabinol and oxandrolone, unless contraindicated or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**MEGACE ES** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

BREAST CANCER AND ENDOMETRIAL CANCER: Megestrol acetate is being used for palliative treatment.

#### **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

#### **Prescriber Restrictions:**

#### **Coverage Duration:**

12 months.

#### Other Criteria:

ANOREXIA AND CACHEXIA ASSOCIATED WITH AIDS: Failure of dronabinol and oxandrolone, unless contraindicated or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**MEKINIST** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

MELANOMA: Positive for BRAF V600E or V600K mutation. NON-SMALL CELL LUNG CANCER, ANAPLASTIC THYROID CANCER: Positive for BRAF V600E mutation.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

NON-SMALL CELL LUNG CANCER, ANAPLASTIC THYROID CANCER: Prescribed in combination with Tafinlar.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**MEKTOVI** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Positive for BRAF V600E or V600K mutation.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

MELANOMA: Prescribed in combination with Braftovi. COLON CANCER, RECTAL CANCER: Prescribed in combination with Braftovi and either Erbitux or Vectibix.

Prior Authorization Group Description:
METAXALONE
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

# **Medicare Part D – 2020**

METHAMPHETAMINE
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Treatment of obesity.
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.

**Prior Authorization Group Description:** 

Prior Authorization Group Description:
METHOCARBAMOL
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

#### Medicare Part D - 2020

# **Prior Authorization Group Description:**

METHOTREXATE INJ

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

**Exclusion Criteria:** 

#### **Required Medical Information:**

**Age Restrictions:** 

#### **Prescriber Restrictions:**

POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS, RHEUMATOID ARTHRITIS: Prescribed by or in consultation with a rheumatologist. PSORIASIS: Prescribed by or in consultation with a rheumatologist or a dermatologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

Failure of generic methotrexate injection, unless contraindicated or clinically significant adverse effects are experienced.

### **Medicare Part D – 2020**

Prior Authorization Group Description:
MIRVASO
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Erythema of rosacea with papules or pustules: Failure of topical metronidazole, oral doxycycline or Finacea, unless

contraindicated or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**MOZOBIL** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Member is scheduled to receive autologous stem cell transplantation.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

Prescribed in combination with a granulocyte-colony stimulating factor (G-CSF) (i.e., Neupogen, Zarxio, Granix, or Nivestym).

#### Medicare Part D - 2020

or Authorization Group Descrip	otion:	

**MULPLETA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Recent (within the past 14 days) platelet count is less than  $50 \times 10^9$ /L. Member is scheduled to undergo a medical or dental procedure within the next 30 days.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a hematologist, hepatologist, or gastroenterologist.

#### **Coverage Duration:**

4 weeks.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**NATPARA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Recent (dated within the last 30 days) serum calcium level is greater than 7.5 mg/dL. Recent (dated within the last 30 days) lab result shows sufficient 25-hydroxyvitamin D stores (at least 50 nmol/L or 20 ng/mL). CONTINUATION OF THERAPY: Maintained on therapy with positive response as evidenced by a recent (dated within the last 90 days) serum calcium level within 8-9 mg/dL.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an endocrinologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

Failure of an active form of vitamin D (e.g., calcitriol) unless contraindicated or clinically significant adverse events are experienced.

### **Medicare Part D – 2020**

Prior	Authorization	Group	<b>Description:</b>	

NAYZILAM

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Member is currently on a stable regimen of antiepileptic drugs.

#### **Age Restrictions:**

Age greater than or equal to 12 years.

#### **Prescriber Restrictions:**

Prescribed by or in consultation with neurologist.

#### **Coverage Duration:**

12 months.

### Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>	

**NERLYNX** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Human epidermal growth factor receptor 2 (HER2)-positive breast cancer.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

#### **Coverage Duration:**

12 months.

Prior Authorization Group Description:
NEULASTA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Mobilization of peripheral-blood progenitor cells prior to autologous transplantation. Supportive care post autologous hematopoietic cell transplantation.
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

<b>Prior Authorization Group Description:</b>		
NEUPOGEN		
Prior Authorization Indication:		
All FDA-approved indications not otherwise excluded from Part D.		
Off Label Uses:		
Myelodysplastic syndrome. Neutropenia in patients with HIV/AIDS.		
Exclusion Criteria:		
Required Medical Information:		
Age Restrictions:		
Prescriber Restrictions:		
Coverage Duration:		
12 months.		
Other Criteria:		

### Medicare Part D - 2020

Prior Authorization Group Description:		

**NINLARO** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

MULTIPLE MYELOMA: Prescribed in combination with dexamethasone.

Prior Authorization Group Description:
NIVESTYM
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Myelodysplastic syndrome. Neutropenia in patients with HIV/AIDS. Hematopoietic syndrome of acute radiation syndrome.
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

Prior Authorization Group Description:
NORTHERA
<b>Prior Authorization Indication:</b>
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
<b>Coverage Duration:</b>
12 months.
Other Criteria:

#### Medicare Part D - 2020

<b>Prior Authorization Group De</b>	escription:
-------------------------------------	-------------

**NUBEQA** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Concurrent use of a gonadotropin-releasing hormone (GnRH) analog (e.g., Zoladex, Vantas, leuprolide/Lupron Depot, Eligard, Trelstar, Firmagon) or past bilateral orchiectomy. Disease is not metastatic.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or urologist.

### Coverage Duration:

12 months.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**NUCALA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

ASTHMA OR EOSINOPHILIC GRANULOMATOSIS WITH POLYANGIITIS: Blood eosinophil count of greater than or equal to 150 cells/mcL within the past 3 months.

#### **Age Restrictions:**

ASTHMA: 6 years of age or older.

#### **Prescriber Restrictions:**

ASTHMA: Prescribed by or in consultation with an allergist, pulmonologist, or immunologist. EOSINOPHILIC GRANULOMATOSIS WITH POLYANGIITIS: Prescribed by or in consultation with a pulmonologist, immunologist, rheumatologist, or nephrologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

ASTHMA: Prescribed in combination with ONE inhaled corticosteroid (e.g., beclomethasone, budesonide, flunisolide, fluticasone, mometasone, ciclesonide), unless contraindicated or clinically significant adverse effects are experienced AND Prescribed in combination with ONE long-acting beta-agonist (e.g., salmeterol, formoterol, vilanterol), unless contraindicated or clinically significant adverse effects are experienced. EOSINOPHILIC GRANULOMATOSIS WITH POLYANGIITIS: Failure of ONE glucocorticoid, unless contraindicated or clinically significant adverse events are experienced.

Prior Authorization Group Description:	
NUEDEXTA	
Prior Authorization Indication:	
All FDA-approved indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	
Required Medical Information:	
Age Restrictions:	
Prescriber Restrictions:	
Prescribed by or in consultation with a neurologist.	
Coverage Duration:	
12 months.	
Other Criteria:	

<b>Prior Authorization Group Description:</b>
NUPLAZID
<b>Prior Authorization Indication:</b>
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
<b>Coverage Duration:</b>
12 months.
Other Criteria:

#### Medicare Part D - 2020

**NUZYRA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Request is for continuation of therapy initiated in an acute care hospital from which member was discharged OR Culture and sensitivity (C&S) report for the current infection shows isolated pathogen is an organism susceptible to omadacycline, unless provider confirms that obtaining a C&S report is not feasible.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

#### **Coverage Duration:**

14 days.

#### Other Criteria:

For members initiating Nuzyra therapy outside of an acute care hospital, one of the following (a, b, or c): a) If a C&S report is available: Failure of 2 antibiotics to which the isolated pathogen is susceptible (if available) per C&S report, unless all are contraindicated or clinically significant adverse effects are experienced. b) C&S report shows resistance or lack of susceptibility of the isolated pathogen to all antibiotics FDA-approved for member's diagnosis. c) If provider confirms that obtaining a C&S report is not feasible: Failure of 2 antibiotics indicated for member's diagnosis (if available), unless all are contraindicated or clinically significant adverse effects are experienced.

# **Medicare Part D – 2020**

**Prior Authorization Group Description:** 

OCALIVA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a hepatologist or gastroenterologist.
Coverage Duration:
12 months.
Other Criteria:
Prescribed in combination with ursodeoxycholic acid unless contraindicated or clinically significant adverse effects are experienced to ursodeoxycholic acid.

#### Medicare Part D - 2020

# **Prior Authorization Group Description: OCREVUS Prior Authorization Indication:** All FDA-approved indications not otherwise excluded from Part D. Off Label Uses: **Exclusion Criteria: Required Medical Information: Age Restrictions: Prescriber Restrictions:** Prescribed by or in consultation with a neurologist. **Coverage Duration:** 12 months. Other Criteria: RELAPSING-REMITTING MULTIPLE SCLEROSIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Aubagio, Tecfidera, Gilenya, Avonex, Betaseron, Plegridy, glatiramer, Copaxone, Glatopa, Extavia or Rebif.

Prior Authorization Group Description:
ODOMZO
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with an oncologist.
Coverage Duration:
12 months.
Other Criteria:

#### Medicare Part D - 2020

Prior	Authorization	Group	<b>Description:</b>

**OFEV** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

SYSTEMIC SCLEROSIS (SSc) ASSOCIATED INTERSTITIAL LUNG DISEASE: Pulmonary fibrosis on high resolution computed tomography (HRCT). Additional signs of SSc are identified (examples may include but are not limited to skin thickening of the fingers, fingertip lesions, telangiectasia, abnormal nailfold capillaries, Raynaud's phenomenon, pulmonary arterial hypertension, SSc-related autoantibodies - anticentromere, anti-topoisomerase I [anti-Scl-70], anti-RNA polymerase III). CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE WITH A PROGRESSIVE PHENOTYPE: For new starts only: confirmation of both of the following within the past 24 months (a and b): a) pulmonary fibrosis affecting more than 10% of lung volume on HRCT and b) confirmation of one of the following (i or ii): i) a relative decline in the forced vital capacity (FVC) of 10% or more of the predicted value, or ii) a relative decline in the FVC of 5% to less than 10% of the predicted value plus either worsening of respiratory symptoms or an increased extent of fibrosis on HRCT.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

SYSTEMIC SCLEROSIS ASSOCIATED INTERSTITIAL LUNG DISEASE, CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE WITH A PROGRESSIVE PHENOTYPE: Prescribed by or in consultation with pulmonologist.

#### **Coverage Duration:**

12 months.

### Medicare Part D – 2020

**Prior Authorization Group Description:** 

OLUMIANT
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a rheumatologist.
Coverage Duration:
12 months.
Other Criteria:
Failure of one of the following agents, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine, or auranofin. Failure

of at least one TNF inhibitor unless contraindicated or clinically significant adverse effects are experienced.

## Medicare Part D – 2020

ONUREG
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
ACUTE MYELOID LEUKEMIA (AML): Prescribed by or in consultation with an oncologist or hematologist.
Coverage Duration:
End of Plan Year.

## Other Criteria:

**Prior Authorization Group Description:** 

AML: Medical justification supports inability to use subcutaneous or intravenous azacitidine (e.g., contraindication to excipients).

Prior Authorization Group Description:	
OPSUMIT	
<b>Prior Authorization Indication:</b>	
All FDA-approved indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	
Required Medical Information:	
Age Restrictions:	
Prescriber Restrictions:	
<b>Coverage Duration:</b>	
12 months.	
Other Criteria:	

Prior Authorization Group Description:
ORENITRAM
<b>Prior Authorization Indication:</b>
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

#### Medicare Part D - 2020

## **Prior Authorization Group Description:**

**ORILISSA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

For 200 mg twice daily requests, members with osteoporosis.

### **Required Medical Information:**

Continuation of therapy: improvement in dysmenorrhea, dyspareunia, pelvic pain/induration/tenderness, or size of endometrial lesions. Total duration of therapy has not exceeded 6 months for 200 mg twice daily or 24 months for 150 mg once daily dosing.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a gynecologist.

### **Coverage Duration:**

200 mg twice daily: 6 months. 150 mg once daily: 12 months.

## Other Criteria:

Failure of ONE non-steroidal anti-inflammatory drug (e.g., ibuprofen, naproxen, fenoprofen, ketoprofen, mefenamic acid, meclofenamate, indomethacin, tolmetin, diclofenac, etodolac, diflunisal, meloxicam, piroxicam) or ONE progestin-containing agent (e.g., norethindrone, ethinyl estradiol with (desogestrel, ethynodiol diacetate, drospirenone, etonogestrel, levonorgestrel, norelgestromin, norethindrone, norgestimate, or norgestrel), estradiol valerate/dienogest, mestranol/norethindrone, depot injectable medroxyprogesterone acetate), unless contraindicated or clinically significant adverse effects are experienced.

## Medicare Part D – 2020

**Prior Authorization Group Description:** 

All FDA-approved indications not otherwise excluded from Part D.

**Prior Authorization Indication:** 

ORKAMBI

Off Label Uses:

**Exclusion Criteria:** 

Required Medical Information:	
Presence of homozygous F508del mutation in an FDA-cleared cystic fibrosis mutation test.	
Age Restrictions:	
2 years of age or older.	
Prescriber Restrictions:	
Coverage Duration:	
12 months.	
Other Criteria:	

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**OXBRYTA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

SICKLE CELL DISEASE: Disease is associated with one of the following genotypes: Homozygous hemoglobin S, Hemoglobin S beta 0-thalassemia, Hemoglobin S beta+ thalassemia, Hemoglobin SC. Member has a hemoglobin level between 5.5 and 10.5 g/dL. Member meets one of the following (a or b): a) Member experienced at least 1 vaso-occlusive crisis (VOC) within the past 6 months while on hydroxyurea, OR b) Member has intolerance or contraindication to hydroxyurea and has experienced at least 1 VOC within the past 12 months. Oxbryta is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced. Oxbryta is not prescribed concurrently with Adakveo. CONTINUATION OF THERAPY, SICKLE CELL DISEASE: Member is responding positively to therapy as evidenced by an increase in Hb level from baseline of at least 1 g/dL. Oxbryta is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced. Oxbryta is not prescribed concurrently with Adakveo.

#### **Age Restrictions:**

SICKLE CELL DISEASE: Age greater than or equal to 12 years.

## **Prescriber Restrictions:**

SICKLE CELL DISEASE: Prescribed by or in consultation with a hematologist.

#### **Coverage Duration:**

6 months.

#### Other Criteria:

SICKLE CELL DISEASE: Failure of L-glutamine, unless contraindicated or clinically significant adverse effects are experienced.

# **Medicare Part D – 2020**

**Prior Authorization Group Description:** 

OXERVATE

<b>Prior Authorization Indication:</b>	
All FDA-approved indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	
Required Medical Information:	
Age Restrictions:	
Prescriber Restrictions:	
Prescribed by or in consultation with an ophthalmologist.	
Coverage Duration:	
12 months.	
Other Criteria:	

#### Medicare Part D - 2020

Prior Authorization G	Group Description:	

**PALYNZIQ** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

## **Exclusion Criteria:**

## **Required Medical Information:**

Recent (within 90 days) phenylalanine (Phe) blood level is greater than 600 micromol/L. CONTINUATION OF THERAPY: Positive response as evidenced by one of the following (a, b, or c): a) Blood Phe level has decreased by at least 20% from pre-treatment baseline, b) Blood Phe level is less than or equal to 600 micromol/L, c) Member has been using 20 mg per day for at least 6 months, but a dose titration to 40 mg per day is being requested after failure to meet therapeutic targets (a or b above).

## **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an endocrinologist, metabolic disease specialist, or genetic disease specialist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

Prior Authorization Group Description:
PEMAZYRE
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.
Coverage Duration:
12 months.
Other Criteria:

Prior Authorization Group Description:
PENNSAID
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of ONE oral non-steroidal anti-inflammatory drug (e.g., ibuprofen, naproxen, fenoprofen, ketoprofen, mefenamic acid, meclofenamate, indomethacin, tolmetin, diclofenac, etodolac, diflunisal, meloxicam, piroxicam) unless all are contraindicated or clinically significant adverse effects are experienced. Failure of either diclofenac 1.5% topical solution or diclofenac 1% topical gel, unless contraindicated or clinically significant adverse effects are experienced.

## Medicare Part D – 2020

Prior Authorization Group Description:
PERSERIS
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Member meets one of the following (a or b): a) therapy was initiated in an inpatient setting during a recent (within 60 days) hospital admission, OR b) failure of TWO of the following atypical antipsychotics, unless contraindicated or clinically significant adverse effects are experienced: aripiprazole, ziprasidone, quetiapine, olanzapine,

risperidone, asenapine, iloperidone, paliperidone.

## Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**PHENOBARBITAL** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

## **Exclusion Criteria:**

#### **Required Medical Information:**

## **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

## **Prescriber Restrictions:**

#### **Coverage Duration:**

12 months.

## Other Criteria:

Partial seizures: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: carbamazepine, phenytoin, topiramate, tiagabine, levetiracetam, gabapentin, lamotrigine, oxcarbazepine, primidone or divalproex. Generalized seizures: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: carbamazepine, phenytoin, topiramate, levetiracetam, primidone or lamotrigine.

## Medicare Part D - 2020

Prior Authorization	Group I	Description:
---------------------	---------	--------------

**PHENTERMINE** 

#### **Prior Authorization Indication:**

Short term, 8 to 12 weeks, adjunct in a regimen of weight reduction based on exercise, behavioral modification and caloric restriction in the management of exogenous obesity for patients with an initial body mass index, BMI, greater than or equal to 30 kg per m^2 or greater than or equal to 27 kg per m^2 in the presence of other risk factors (e.g., hypertension, diabetes, dyslipidemia).

## Off Label Uses:

## **Exclusion Criteria:**

## **Required Medical Information:**

BMI is greater than or equal to 30 kg/m<sup>2</sup> OR BMI is greater than or equal to 27 kg/m<sup>2</sup> with one or more of the following severe co-morbid conditions 1. Coronary artery/heart disease 2. Diabetes 3. Dyslipidemia 4. Hypertension 5. Obstructive sleep apnea.

### **Age Restrictions:**

#### **Prescriber Restrictions:**

## **Coverage Duration:**

Three months.

#### Other Criteria:

## Medicare Part D - 2020

## **Prior Authorization Group Description:**

**PIQRAY** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

## **Exclusion Criteria:**

## **Required Medical Information:**

Hormone receptor (HR)-positive (i.e., estrogen or progesterone receptor [ER/PR]-positive), HER2-negative, advanced (locally recurrent) or metastatic, and positive for PIK3CA-mutation.

#### **Age Restrictions:**

## **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

## **Coverage Duration:**

12 months.

## Other Criteria:

Prescribed in combination with fulvestrant after disease progression on or after an endocrine therapy (e.g., anastrozole, exemestane, fulvestrant, toremifene, letrozole, tamoxifen, or megestrol acetate).

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**PRALUENT** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

GENETICALLY MEDIATED PRIMARY HYPERLIPIDEMIA (INCLUDING HETEROZYGOUS FAMILIAL HYPERCHOLESTEROLEMIA): Confirmation (e.g., medical records, chart notes, laboratory values) of LDL level suggestive of a diagnosis of familial hypercholesterolemia (e.g., adults: LDL of 190 mg/dL or greater). NON-GENETICALLY MEDIATED PRIMARY HYPERLIPIDEMIA: Request meets both of the following (a and b): a) Confirmation of a LDL of 100 mg/dL or greater AND b) a diagnosis of secondary hyperlipidemia has been ruled out with confirmation of absence of all of the following potential causes of elevated cholesterol (a-e): a) hypothyroidism, b) obstructive liver disease, c) renal disease, d) nephrosis, e) medications which can increase lipid levels including, but not limited to: glucocorticoids, sex hormones, antipsychotics, antiretrovirals, immunosuppressive agents, retinoic acid derivatives. HYPERCHOLESTEROLEMIA WITH HISTORY OF CLINICAL ATHEROSCLEROTIC CARDIOVASCULAR DISEASE (ASCVD): Confirmation of an LDL of 70 mg/dL or greater AND history of clinical ASCVD defined as one of the following: Acute coronary syndromes, Myocardial infarction, Stable or unstable angina, Coronary or other arterial revascularization (e.g., percutaneous coronary intervention or coronary artery bypass graft surgery), Stroke, Peripheral artery disease presumed to be of atherosclerotic origin, Transient ischemic attack (TIA), Clinically significant coronary heart disease (CHD) diagnosed by invasive or noninvasive testing (such as coronary angiography, stress test using treadmill, stress echocardiography, or nuclear imaging), Carotid artery occlusion greater than 50% without symptoms, Renal artery stenosis or renal artery stent procedure. CONTINUATION OF THERAPY: Confirmation of LDL reduction while on Praluent therapy AND, if tolerated, confirmation of continued statin therapy at the maximally tolerated dose.

## Age Restrictions:

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist.

## **Coverage Duration:**

6 months.

#### Other Criteria:

Failure of two of the following at maximally tolerated doses, unless contraindicated or clinically significant adverse effects are experienced: atorvastatin, rosuvastatin, simvastatin, ezetimibe/simvastatin, pitavastatin, pravastatin, fluvastatin, or lovastatin.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**PRETOMANID** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

## **Exclusion Criteria:**

#### **Required Medical Information:**

Confirmed resistance to fluoroquinolones, unless contraindicated or clinically significant adverse effects are experienced. CONTINUATION OF THERAPY: Confirmation of delayed culture conversion and total duration of pretomanid therapy has not exceeded 9 months.

#### **Age Restrictions:**

## **Prescriber Restrictions:**

Prescribed by or in consultation with an expert in the treatment of tuberculosis.

### **Coverage Duration:**

Initial: 6 months. Reauthorization: 3 months.

#### Other Criteria:

Prescribed in combination with bedaquiline and linezolid. CONTINUATION OF THERAPY: Member meets one of the following (a or b): a) Prescribed in combination with bedaquiline and linezolid OR b) If member has completed at least 4 weeks of linezolid therapy, member continues to receive pretomanid in combination with bedaquiline.

## Medicare Part D - 2020

## **Prior Authorization Group Description:**

**PREVYMIS** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

## **Exclusion Criteria:**

Member is receiving pimozide or ergot alkaloids. Member is receiving cyclosporine co-administered with pitavastatin or simvastatin.

## **Required Medical Information:**

#### **Age Restrictions:**

## **Prescriber Restrictions:**

Prescribed by or in consultation with an oncology, hematology, infectious disease, or transplant specialist.

#### **Coverage Duration:**

Through day 100 post-transplantation.

## Other Criteria:

Failure of generic valacyclovir or generic ganciclovir, unless contraindicated, clinically significant adverse effects are experienced, or member is at high risk for CMV.

## Medicare Part D – 2020

**Prior Authorization Group Description:** 

PROLASTIN C			
<b>Prior Authorization Indication:</b>			
All FDA-approved indications not other	wise excluded from	n Part D.	
Off Label Uses:			
<b>Exclusion Criteria:</b>			
Required Medical Information:			
Age Restrictions:			
Prescriber Restrictions:			
Prescribed by or in consultation with a	oulmonologist.		
<b>Coverage Duration:</b>			
12 months.			
Other Criteria:			

## Medicare Part D – 2020

**Prior Authorization Group Description:** 

(Femara)].

PROLIA
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
PROSTATE CANCER: Receiving or has received androgen deprivation therapy [e.g., leuprolide (Lupron),

bicalutamide (Casodex) or Nilandron]. BREAST CANCER: Receiving or has received adjuvant endocrine therapy [e.g., tamoxifen or aromatase inhibitors such as anastrozole (Arimidex), exemestane (Aromasin) or letrozole

## Medicare Part D - 2020

## **Prior Authorization Group Description:**

**PROMACTA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### **Off Label Uses:**

## **Exclusion Criteria:**

#### **Required Medical Information:**

Thrombocytopenia in Chronic Hepatitis C: Confirmation of current or planned interferon-based treatment of chronic hepatitis C.

#### **Age Restrictions:**

### **Prescriber Restrictions:**

CHRONIC IMMUNE THROMBOCYTOPENIA, SEVERE APLASTIC ANEMIA: Prescribed by or in consultation with a hematologist. THROMBOCYTOPENIA IN CHRONIC HEPATITIS C: Prescribed by or in consultation with a hematologist, gastroenterologist, or an infectious disease specialist.

## **Coverage Duration:**

12 months.

## Other Criteria:

Chronic Immune (Idiopathic) Thrombocytopenia: Failure of a corticosteroid (e.g., prednisone, methylprednisolone, or dexamethasone), unless contraindicated or clinically significant adverse effects are experienced.

## Medicare Part D - 2020

|--|

## **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

**PROTOPIC** 

**Exclusion Criteria:** 

## **Required Medical Information:**

## **Age Restrictions:**

Tacrolimus 0.1%: 16 years and older.

## **Prescriber Restrictions:**

## **Coverage Duration:**

12 months.

#### Other Criteria:

Failure of two medium to high potency topical corticosteroids (e.g., amcinonide, fluticasone propionate, triamcinolone acetonide, betamethasone valerate, fluocinolone acetonide, hydrocortisone butyrate, mometasone furoate, desoximetasone, fluocinonide or betamethasone dipropionate), unless contraindicated or clinically significant adverse effects are experienced.

## Medicare Part D – 2020

<b>Prior Authorization Indication:</b>
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Multiple sclerosis-related fatigue.
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

**Prior Authorization Group Description:** 

PROVIGIL

## Medicare Part D - 2020

**Prior Authorization Group Description:** 

**PURIXAN** 

Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with oncologist or hematologist.
Coverage Duration:
12 months.
Other Criteria:
One of the following: Failure of mercaptopurine tablets, unless contraindicated or clinically significant adverse

effects are experienced OR member has a swallowing disorder or an inability to swallow tablets or capsules..

## Medicare Part D – 2020

**Prior Authorization Group Description:** 

QINLOCK
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.
Coverage Duration:
12 months.
Other Criteria:
GASTROINTESTINAL STROMAL TUMOR: For members with PDGFRA exon 18 mutation, failure of Ayvakit, unless contraindicated or clinically significant adverse effects are experienced.

## Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**QSYMIA** 

#### **Prior Authorization Indication:**

As an adjunct to a reduced-calorie diet and increased physical activity for chronic weight management in adults with an initial body mass index (BMI) of 30 kg/m<sup>2</sup> or greater (obese) or 27 kg/m<sup>2</sup> or greater (overweight) in the presence of at least one weight-related comorbidity such as hypertension, type 2 diabetes mellitus, or dyslipidemia.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

BMI is greater than or equal to 30 kg/m<sup>2</sup> OR BMI is greater than or equal to 27 kg/m<sup>2</sup> with one or more of the following severe co-morbid conditions 1. Coronary artery/heart disease 2. Diabetes 3. Dyslipidemia 4. Hypertension 5. Obstructive sleep apnea. CONTINUATION OF THERAPY: If 3% weight loss is not achieved after 12 weeks on 7.5 mg/46 mg, then dose must be escalated or drug discontinued. If dose is escalated, an additional 12 weeks will be approved. Reauthorization every 6 months for the first year will require confirmation of at least 5% weight loss from baseline body weight. Therapy beyond the first year can be authorized every 6 months with confirmation of weight maintenance.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

#### **Coverage Duration:**

Initial: 12 weeks. First reauthorization: 12 weeks. Subsequent reauthorizations: 6 months.

### Other Criteria:

## Medicare Part D - 2020

## **Prior Authorization Group Description:**

QUALAQUIN

## **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

## Off Label Uses:

Babesiosis. Plasmodium vivax malaria.

## **Exclusion Criteria:**

For the treatment or prevention of nocturnal leg cramps.

## **Required Medical Information:**

## **Age Restrictions:**

## **Prescriber Restrictions:**

## **Coverage Duration:**

Malaria: 7 days. Babesiosis: 7-10 days.

## Other Criteria:

Plasmodium vivax malaria: Infection is chloroquine-resistant.

## Medicare Part D - 2020

## **Prior Authorization Group Description:**

**RADICAVA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

## **Exclusion Criteria:**

#### **Required Medical Information:**

Forced vital capacity greater than or equal to 80%, disease duration of less than or equal to 2 years, functionally retains most activities of daily living (defined as a baseline revised ALS Functional Rating Scale (ALSFRS-R) score with greater than or equal to 2 points in each of the 12 items, meets diagnostic criteria of definite or probable amyotrophic lateral sclerosis (ALS) based on El Escorial revised criteria. CONTINUATION OF THERAPY: Member continues to retain most activities of daily living, forced vital capacity greater than or equal to 80%, and ALSFRS-R score with greater than or equal to 2 points in each of the 12 items.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist.

## **Coverage Duration:**

6 months.

#### Other Criteria:

Prescribed in combination with riluzole unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:
RAYALDEE
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Patient has stage 3 or 4 chronic kidney disease (CKD) and serum total 25-hydroxyvitamin D level less than 30 $$ ng/mL.
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

REBLOZYL

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

TRANSFUSION DEPENDENT BETA-THALASSEMIA: Total volume of transfusions exceeds 6 red blood cell units within the last 6 months. No transfusion free period for greater than or equal to 35 days within the last 6 months. CONTINUATION OF THERAPY, TRANSFUSION DEPENDENT BETA THALASSEMIA: Member meets one of the following (a or b): a) Member is responding positively to therapy as evidenced by at least a 33% reduction in transfusion burden from baseline, b) Request is for a dose increase. MYELODYSPLASTIC SYNDROMES (MDS): Member requires 2 or more RBC units per 8 weeks confirmed for at least the last 16 weeks. Member has either a ring sideroblast of at least 15% of erythroid precursors in bone marrow or ring sideroblast of at least 5% if SF3B1 mutation is present. Member does not have del(5q) cytogenetic abnormality. CONTINUATION OF THERAPY, MDS: Member meets one of the following (a or b): a) Member is responding positively to therapy as evidenced by a decreased transfusion burden, b) Request is for a dose increase.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

TRANSFUSION DEPENDENT BETA-THALASSEMIA: Prescribed by or in consultation with a hematologist. MDS: Prescribed by or in consultation with a hematologist or oncologist.

#### **Coverage Duration:**

TRANSFUSION DEPENDENT BETA-THALASSEMIA, MDS: Initial: 2 months. Reauthorization: 6 months.

#### Other Criteria:

MDS: Failure of an erythropoiesis-stimulating agent used in combination with a granulocyte colony stimulating factor, unless clinically significant adverse effects are experienced, all are contraindicated, or confirmation of current serum erythropoietin greater than 500 mU/mL.

#### Medicare Part D - 2020

## **Prior Authorization Group Description:**

**REMICADE** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

Wegener's Granulomatosis.

## **Exclusion Criteria:**

#### **Required Medical Information:**

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Psoriatic Arthritis/Plaque Psoriasis: Prescribed by or in consultation with a rheumatologist or dermatologist. Crohn's Disease/Ulcerative Colitis: Prescribed by or in consultation with a gastroenterologist. Rheumatoid Arthritis/Ankylosing Spondylitis: Prescribed by or in consultation with a rheumatologist.

### **Coverage Duration:**

12 months.

#### Other Criteria:

Rheumatoid Arthritis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. Plaque Psoriasis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin.

#### Medicare Part D - 2020

## Prior Authorization Group Description:

**RENFLEXIS** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

**Exclusion Criteria:** 

#### **Required Medical Information:**

## **Age Restrictions:**

#### **Prescriber Restrictions:**

Psoriatic Arthritis/Plaque Psoriasis: Prescribed by or in consultation with a rheumatologist or dermatologist. Crohn's Disease/Ulcerative Colitis: Prescribed by or in consultation with a gastroenterologist. Rheumatoid Arthritis/Ankylosing Spondylitis: Prescribed by or in consultation with a rheumatologist.

#### **Coverage Duration:**

12 months.

## Other Criteria:

Rheumatoid Arthritis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. Plaque Psoriasis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin.

#### Medicare Part D - 2020

## **Prior Authorization Group Description:**

REPATHA

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

## **Required Medical Information:**

GENETICALLY MEDIATED PRIMARY HYPERLIPIDEMIA (INCLUDING HETEROZYGOUS OR HOMOZYGOUS FAMILIAL HYPERCHOLESTEROLEMIA): Confirmation (e.g., medical records, chart notes, laboratory values) of LDL level suggestive of a diagnosis of familial hypercholesterolemia (e.g., adults: LDL 190 mg/dL or greater). NON-GENETICALLY MEDIATED PRIMARY HYPERLIPIDEMIA: Request meets both of the following (a and b): a) Confirmation of a LDL of 100 mg/dL or greater AND b) a diagnosis of secondary hyperlipidemia has been ruled out with confirmation of absence of all of the following potential causes of elevated cholesterol (a-e): a) hypothyroidism, b) obstructive liver disease, c) renal disease, d) nephrosis, e) medications which can increase lipid levels including, but not limited to: glucocorticoids, sex hormones, antipsychotics, antiretrovirals, immunosuppressive agents, retinoic acid derivatives. HYPERCHOLESTEROLEMIA WITH HISTORY OF CLINICAL ATHEROSCLEROTIC CARDIOVASCULAR DISEASE (ASCVD): Confirmation of an LDL of 70 mg/dL or greater AND history of clinical ASCVD defined as one of the following: Acute coronary syndromes, Myocardial infarction, Stable or unstable angina, Coronary or other arterial revascularization (e.g., percutaneous coronary intervention or coronary artery bypass graft surgery), Stroke, Peripheral artery disease presumed to be of atherosclerotic origin, Transient ischemic attack (TIA), Clinically significant coronary heart disease (CHD) diagnosed by invasive or noninvasive testing (such as coronary angiography, stress test using treadmill, stress echocardiography, or nuclear imaging), Carotid artery occlusion greater than 50% without symptoms, Renal artery stenosis or renal artery stent procedure. CONTINUATION OF THERAPY: Confirmation of LDL reduction while on Repatha therapy AND, if tolerated, confirmation of continued statin therapy at the maximally tolerated dose.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist.

#### **Coverage Duration:**

6 months

### Other Criteria:

Failure of two of the following at maximally tolerated doses, unless contraindicated or clinically significant adverse effects are experienced: atorvastatin, rosuvastatin, simvastatin, ezetimibe/simvastatin, pitavastatin, pravastatin, fluvastatin, or lovastatin.

Prior Authorization Group Description:
RETEVMO
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.
Coverage Duration:
12 months.
Other Criteria:

Prior Authorization Group Description:
REVATIO
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Members on concomitant nitrates (e.g., Nitrodur, Nitrobid, Nitrostat, Isordil, Ismo). Members on concomitant guanylate cyclase stimulator, such as riociguat (Adempas).
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

<b>Prior Authorization Group Description:</b>
REVCOVI
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with an immunologist.
<b>Coverage Duration:</b>
12 months.
Other Criteria:

<b>Prior Authorization Group Description:</b>
REVLIMID
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with an oncologist or hematologist.
Coverage Duration:
12 months.
Other Criteria:

Prior Authorization Group Description:
REXULTI
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of aripiprazole and one of the following generic atypical antipsychotics, unless contraindicated or clinically significant adverse effects are experienced: risperidone, olanzapine, quetiapine, ziprasidone.

Prior Authorization Group Description:
RINVOQ
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a rheumatologist.
Coverage Duration:
12 months.
Other Criteria:
RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin.

# Medicare Part D - 2020

**ROZLYTREK** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

ROS1-POSITIVE NON-SMALL CELL LUNG CANCER: Confirmation of a ROS1 mutation. Member has not received prior ROS1 targeted therapy (e.g., Xalkori, Zykadia, Lorbrena). NTRK FUSION-POSITIVE SOLID TUMOR: Confirmation of an NTRK gene fusion (e.g., ETV6-NTRK3, TPM3-NTRK1). Member has not received prior NTRK targeted therapy (e.g., Vitrakvi).

# **Age Restrictions:**

NTRK FUSION-POSITIVE SOLID TUMOR: Age greater than or equal to 12 years.

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# Medicare Part D - 2020

# **Prior Authorization Group Description:**

**RUBRACA** 

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

OVARIAN CANCER: Mutations in the BRCA genes OR member has a complete or partial response to two or more platinum-based chemotherapy regimens.

#### **Age Restrictions:**

# **Prescriber Restrictions:**

PROSTATE CANCER: Prescribed by or in consultation with an oncologist or urologist. ALL OTHER ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# **Medicare Part D - 2020**

RUZURGI	
<b>Prior Authorization Indication:</b>	
All FDA-approved indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	

#### Required Medical Information:

Confirmation of a baseline clinical muscle strength assessment (examples may include but are not limited to the Quantitative Myasthenia Gravis (QMG) score, triple-timed up-and-go test (3TUG), Timed 25-foot Walk test (T25FW)). CONTINUATION OF THERAPY: Member is responding positively to therapy as evidenced by clinical muscle strength assessments.

# **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist.

# **Coverage Duration:**

12 months.

#### Medicare Part D - 2020

# **Prior Authorization Group Description:**

**RYDAPT** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

Acute Myeloid Leukemia: Positive for the FLT3 mutation.

# **Age Restrictions:**

# **Prescriber Restrictions:**

Acute Myeloid Leukemia: Prescribed by or in consultation with an oncologist or hematologist. Advanced Systemic Mastocytosis: Prescribed by or in consultation with an oncologist, allergist, or immunologist.

#### **Coverage Duration:**

12 months.

# Other Criteria:

Acute Myeloid Leukemia: for induction therapy, prescribed in combination with cytarabine and daunorubicin OR for consolidation therapy, prescribed in combination with cytarabine.

Prior Authorization Group Description:
SECUADO
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Medical justification supports inability to use Saphris (asenapine sublingual tablets) (e.g., contraindications to excipients).

# Medicare Part D – 2020

**Prior Authorization Group Description:** 

methotrexate, cyclosporine or acitretin.

SILIQ

Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a rheumatologist or dermatologist.
<b>Coverage Duration:</b>
12 months.
Other Criteria:
Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced:

# Medicare Part D - 2020

#### **Prior Authorization Group Description:**

SIMPONI(prefilled syringe, auto-injector)

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

# **Exclusion Criteria:**

#### **Required Medical Information:**

# **Age Restrictions:**

#### **Prescriber Restrictions:**

PSORIATIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist or dermatologist. ULCERATIVE COLITIS: Prescribed by or in consultation with a gastroenterologist. RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS: Prescribed by or in consultation with a rheumatologist.

#### **Coverage Duration:**

12 months.

# Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin.

# Medicare Part D - 2020

# **Prior Authorization Group Description:**

SIMPONI ARIA

# **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

**Exclusion Criteria:** 

# **Required Medical Information:**

**Age Restrictions:** 

#### **Prescriber Restrictions:**

PSORIATIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist or dermatologist. RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS: Prescribed by or in consultation with a rheumatologist.

# **Coverage Duration:**

12 months.

# Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin.

# Medicare Part D – 2020

**Prior Authorization Group Description:** 

methotrexate, cyclosporine, or acitretin.

SKYRIZI

Prior Authorization Indication:	
All FDA-approved indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	
Required Medical Information:	
Age Restrictions:	
Prescriber Restrictions:	
Prescribed by or in consultation with a dermatologist or rheumatologist.	
Coverage Duration:	
12 months.	
Other Criteria:	
Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced	

Prior Authorization Group Description:
SOMA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

Prior Authorization Group Description:	
SOMAVERT	
Prior Authorization Indication:	
All FDA-approved indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	
Required Medical Information:	
Age Restrictions:	
Prescriber Restrictions:	
Prescribed by or in consultation with an endocrinologist.	
Coverage Duration:	
12 months.	
Other Criteria:	
Inadequate response to surgery or radiation therapy, unless surgery or radiation therapy is not appropriate for patient.	the

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**SOVALDI** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

Criteria will be applied consistent with current AASLD-IDSA guidance.

#### **Exclusion Criteria:**

### **Required Medical Information:**

Refer to the document, "Recommendations for Testing, Managing, and Treating Hepatitis C," by AASLD-IDSA available at http://www.hcvguidelines.org for drug regimen and duration of treatment based on genotype, treatment status, previous drug regimens used, past medical history and comorbidities.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a gastroenterologist, hepatologist or infectious disease physician.

### **Coverage Duration:**

Criteria will be applied consistent with current AASLD-IDSA guidance.

# Other Criteria:

Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: Mavyret, Harvoni, Epclusa, Vosevi, and Zepatier for applicable genotypes.

#### Medicare Part D - 2020

# **Prior Authorization Group Description:**

**SPRAVATO** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

Currently on an oral antidepressant (must not be an agent previously tried and failed). CONTINUATION OF THERAPY: Member is responding positively to therapy and is using Spravato in combination with an oral antidepressant.

# **Age Restrictions:**

# **Prescriber Restrictions:**

#### **Coverage Duration:**

Initial: 4 weeks. Reauthorization: 6 months.

# Other Criteria:

Failure of two antidepressants (e.g., selective serotonin reuptake inhibitor [SSRI], serotonin-norepinephrine reuptake inhibitor [SNRI], tricyclic antidepressant [TCA], bupropion, mirtazapine) from two different classes, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:
SPRITAM
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Medical justification must be provided why patient cannot take generic levetiracetam tablets or liquid.

#### Medicare Part D - 2020

# **Prior Authorization Group Description:**

**SPRYCEL** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

CHRONIC MYELOGENOUS LEUKEMIA, ACUTE LYMPHOBLASTIC LEUKEMIA: Confirmation that the member has Philadelphia chromosome positive disease.

#### **Age Restrictions:**

# **Prescriber Restrictions:**

CHRONIC MYELOGENOUS LEUKEMIA, ACUTE LYMPHOBLASTIC LEUKEMIA: Prescribed by or in consultation with an oncologist or hematologist. ALL OTHER COVERED ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# Other Criteria:

GASTROINTESTINAL STROMAL TUMOR: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: imatinib, Sutent or Stivarga.

# Medicare Part D – 2020

STELARA IV
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a gastroenterologist.
Coverage Duration:
4 weeks.
Other Criteria:
CROHN'S DISEASE: Failure of Humira or Remicade and one of the following, unless contraindicated or clinical significant adverse effects are experienced: 6-mercaptopurine, azathioprine or methotrexate.

#### Medicare Part D - 2020

# **Prior Authorization Group Description:**

STELARA SC

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

**Exclusion Criteria:** 

# **Required Medical Information:**

# **Age Restrictions:**

#### **Prescriber Restrictions:**

PSORIATIC ARTHRITIS, PLAQUE PSORIASIS: Prescribed by or in consultation with a rheumatologist or dermatologist. CROHN'S DISEASE, ULCERATIVE COLITIS: Prescribed by or in consultation with a gastroenterologist.

#### **Coverage Duration:**

12 months.

# Other Criteria:

PLAQUE PSORIASIS: Failure of ONE of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine, or acitretin. PSORIATIC ARTHRITIS: Failure of one TNF inhibitor (e.g., Humira, Enbrel, Cimzia, Simponi, Simponi Aria, Remicade, Inflectra), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:
STRENSIQ
<b>Prior Authorization Indication:</b>
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
<b>Prescriber Restrictions:</b>
<b>Coverage Duration:</b>
12 months.
Other Criteria:

Prior Authorization Group Description:	
SUBSYS	
Prior Authorization Indication:	
All FDA-approved indications not otherwise excluded from Part D.	
Off Label Uses:	
Exclusion Criteria:	
Required Medical Information:	
Age Restrictions:	
Age 18 or greater.	
Prescriber Restrictions:	
Coverage Duration:	
Through the end of the Plan contract year.	
Other Criteria:	
Patient is already taking and is tolerant to around-the-clock opioid therapy. Patients are considered opioid toler when taking another opioid daily for a week or longer (for example, at least 60 mg of oral morphine per day or equianalgesic dose of another opioid).	

Prior Authorization Group Description:
SUNOSI
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
NARCOLEPSY: Prescribed by or in consultation with a neurologist.
Coverage Duration:
12 months.
Other Criteria:
Failure of armodafinil (Nuvigil) or modafinil (Provigil), unless contraindicated or clinically significant side effects are experienced.

# Medicare Part D - 2020

# **Prior Authorization Group Description:**

**SURMONTIL** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

# Off Label Uses:

Irritable bowel syndrome.

# **Exclusion Criteria:**

# **Required Medical Information:**

# **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

# **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

# Other Criteria:

Depression: Failure of one of the following generic antidepressants, unless contraindicated or clinically significant adverse effects are experienced: bupropion, bupropion SR, bupropion XL, citalopram, desvenlafaxine succinate, duloxetine, escitalopram, fluoxetine, mirtazapine, sertraline, venlafaxine, or venlafaxine XR.

# **Medicare Part D – 2020**

SYMDEKO
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Presence of homozygous F508del mutation or at least one mutation in the CFTR gene that is responsive to tezacaftor/ivacaftor.
Age Restrictions:
Age greater than or equal to 6 years.
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D – 2020

SYMLINPEN
<b>Prior Authorization Indication:</b>
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
<b>Coverage Duration:</b>
12 months.
Other Criteria:
Previous use of mealtime insulin therapy or an insulin pump.

# Medicare Part D – 2020

SYMPAZAN
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a neurologist.
Coverage Duration:
12 months.
Other Criteria:
Medical justification supports inability to use clobazam tablets and oral suspension (e.g., contraindications excipients).

# Medicare Part D - 2020

Prior	Authorizatio	n Group	<b>Description:</b>

**TABRECTA** 

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

NON-SMALL CELL LUNG CANCER: Disease is epidermal growth factor receptor (EGFR) wild-type and anaplastic lymphoma kinase (ALK) negative.

# **Age Restrictions:**

# **Prescriber Restrictions:**

ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

#### Medicare Part D - 2020

Prior Authorization	Group	<b>Description:</b>
---------------------	-------	---------------------

**TAFAMIDIS** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

#### **Exclusion Criteria:**

# **Required Medical Information:**

TRANSTHYRETIN AMYLOID CARDIOMYOPATHY (ATTR-CM): Member has not had a liver transplant. Diagnosis is supported by (a or b): a) Confirmation of amyloid deposition on biopsy and either transthyretin (TTR) precursor protein (e.g., by immunohistochemistry, scintigraphy, mass spectrometry) or a TTR mutation by genetic testing. b) Member meets all of the following (i, ii, and iii): i) Echo, CMR, or PET findings are consistent with cardiac amyloidosis, AND ii) Cardiac update is Grade 2 or 3 on a radionuclide scan utilizing one of the following radiotracers (1, 2, or 3): 1) 99m technetium (Tc)-labeled 3,3-diphosphono-1,2-propanodicarboxylic acid (DPD), 2) 99mTc-labeled pyrophosphate (PYP), or 3) 99mTc-labeled hydroxymethylene diphosphonate (HMDP), AND iii) Each of the following laboratory tests is negative for monoclonal protein (1, 2, and 3): 1) Serum kappa/lambda free light chain ratio analysis, 2) Serum protein immunofixation, 3) Urine protein immunofixation. CONTINUATION OF THERAPY, ATTR-CM: Maintained on therapy with positive response, including but not limited to, improvement or stabilization in any of the following parameters: 1) walking ability, 2) nutrition (e.g., body mass index), 3) cardiac related hospitalization, 4) cardiac procedures or laboratory tests (e.g., Holter monitoring, echocardiography, electrocardiogram, plasma BNP or NT-proBNP, serum troponin).

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a cardiologist.

#### **Coverage Duration:**

12 months.

# Medicare Part D - 2020

**TAGRISSO** 

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

NON-SMALL CELL LUNG CANCER: Disease is positive for either of the following (a or b): a) sensitizing EGFR mutation (e.g., exon 19 deletion or insertion, exon 21 point mutation (L858R, L861Q), exon 18 point mutation (G719X), exon 20 point mutation (S768I)), OR b) T790M mutation.

# **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# Medicare Part D – 2020

All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Age greater than or equal to 12 years.
Prescriber Restrictions:
Prescribed by or in consultation with an immunologist, allergist, hematologist, or rheumatologist
<b>Coverage Duration:</b>
12 months.

**Prior Authorization Group Description:** 

**Prior Authorization Indication:** 

TAKHZYRO

# Medicare Part D – 2020

**Prior Authorization Group Description:** 

TALZENNA
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Confirmation of human epidermal growth factor receptor 2 (HER2)-negative disease and mutation in the BRC genes.
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with an oncologist.
Coverage Duration:
12 months.

# Medicare Part D - 2020

# **Prior Authorization Group Description:**

**TARCEVA** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

NON-SMALL CELL LUNG CANCER: Disease is positive for a sensitizing EGFR mutation (e.g., exon 19 deletion or insertion, exon 21 point mutation (L858R, L861Q), exon 18 point mutation (G719X), exon 20 point mutation (S768I)). RENAL CELL CARCINOMA: Confirmation of non-clear cell histology.

# **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

#### Other Criteria:

PANCREATIC CANCER: Prescribed in combination with gemcitabine.

# Medicare Part D - 2020

TARGRETIN GEL

# **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D. Some medically-accepted indications.

# Off Label Uses:

Primary cutaneous B-cell lymphoma types: primary cutaneous marginal zone lymphoma and primary cutaneous follicle center lymphoma.

# **Exclusion Criteria:**

# **Required Medical Information:**

# **Age Restrictions:**

# **Prescriber Restrictions:**

ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

#### Medicare Part D - 2020

# **Prior Authorization Group Description:**

**TASIGNA** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

CHRONIC MYELOGENOUS LEUKEMIA, ACUTE LYMPHOBLASTIC LEUKEMIA: Confirmation that the member has Philadelphia chromosome positive disease.

#### **Age Restrictions:**

# **Prescriber Restrictions:**

CHRONIC MYELOGENOUS LEUKEMIA, ACUTE LYMPHOBLASTIC LEUKEMIA: Prescribed by or in consultation with an oncologist or hematologist. GASTROINTESTINAL STROMAL TUMOR: Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# Other Criteria:

GASTROINTESTINAL STROMAL TUMOR: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: imatinib, Sutent or Stivarga.

<b>Prior Authorization Group Description:</b>
TAVALISSE
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of a corticosteroid (e.g., prednisone, methylprednisolone, or dexamethasone), unless contraindicated or clinically significant adverse effects are experienced.

# Medicare Part D - 2020

# **Prior Authorization Group Description:**

**TAZVERIK** 

### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

EPITHELIOID SARCOMA: Tumor demonstrates loss of INI1 expression through inactivation, deletion, or mutation of the INI1 (SMARCB-1) gene.

# **Age Restrictions:**

EPITHELIOID SARCOMA: Age 16 years or older.

# **Prescriber Restrictions:**

FOLLICULAR LYMPHOMA: Prescribed by or in consultation with an oncologist or hematologist. EPITHELIOID SARCOMA: Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

### Medicare Part D - 2020

## **Prior Authorization Group Description:**

**TECENTRIQ** 

### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

### Off Label Uses:

# **Exclusion Criteria:**

### **Required Medical Information:**

EXTENSIVE-STAGE SMALL CELL LUNG CANCER: Prescribed in combination with carboplatin and etoposide. TRIPLE NEGATIVE BREAST CANCER: Hormone-receptor (HR)-negative, estrogen-receptor (ER)-negative, and human epidermal growth factor receptor 2 (HER2)-negative disease. Prescribed in combination with protein-bound paclitaxel (nab-paclitaxel). Tumor expresses PD-L1.

### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

### **Coverage Duration:**

12 months.

#### Other Criteria:

NON-SMALL CELL LUNG CANCER: If a known epidermal growth factor receptor (EGFR) or anaplastic lymphoma kinase (ALK) genomic tumor aberration exists, then for ALK+ disease: prior trial of Xalkori, Alecensa, or Zykadia OR for EGFR+ disease: prior trial of Tarceva, Gilotrif, or Iressa.

# Medicare Part D – 2020

**Prior Authorization Group Description:** 

TECFIDERA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a neurologist.
Coverage Duration:
12 months.
Other Criteria:

### Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>
---

**TEGSEDI** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

# **Exclusion Criteria:**

### **Required Medical Information:**

Confirmation of transthyretin (TTR) mutation. Confirmation of amyloid deposition on biopsy or medical justification is provided as to why treatment should be initiated in the presence of a negative biopsy or no biopsy. Member has not had a liver transplant. CONTINUATION OF THERAPY: Maintained on therapy with positive response, including but not limited to improvement in any of the following parameters: 1) neuropathy (motor function, sensation, reflexes, walking ability), 2) nutrition (body mass index), 3) cardiac parameters (Holter monitoring, echocardiography, electrocardiogram, plasma BNP or NT-proBNP, serum troponin), 4) renal parameters (creatinine clearance, urine albumin), 5) ophthalmic parameters (eye exam).

### **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist.

### **Coverage Duration:**

12 months.

### Medicare Part D - 2020

# **Prior Authorization Group Description:**

**TENEX** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

### **Exclusion Criteria:**

# **Required Medical Information:**

# **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

### **Prescriber Restrictions:**

### **Coverage Duration:**

12 months.

# Other Criteria:

Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: amlodipine/benazepril, benazepril/hydrochlorothiazide, captopril, captopril/hydrochlorothiazide, fosinopril, fosinopril/hydrochlorothiazide, lisinopril, lisinopril/hydrochlorothiazide, quinapril, quinapril/hydrochlorothiazide, losartan, losartan/hydrochlorothiazide, valsartan, valsartan/hydrochlorothiazide, irbesartan, irbesartan/hydrochlorothiazide, candesartan, candesartan/hydrochlorothiazide, carvedilol, labetalol, acebutolol, atenolol, bisoprolol/hydrochlorothiazide, timolol, nadolol, propranolol, metoprolol, metoprolol/hydrochlorothiazide, pindolol, nifedipine SR, amlodipine, nicardipine.

#### Medicare Part D - 2020

### **Prior Authorization Group Description:**

**TEPEZZA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

### **Required Medical Information:**

THYROID EYE DISEASE (TED): Member has active TED with a clinical activity score (CAS) of at least 4 or greater. Member is euthyroid with confirmation of a recent (within the last 30 days) free thyroxine (FT4) and free triiodothyronine (FT3) levels within the laboratory defined reference range. Member has not had previous surgical intervention for TED. Member does not require surgical ophthalmological intervention. Member has not received 8 or more infusions lifetime (including the initial 10 mg/kg first infusion). CONTINUATION OF THERAPY, TED: Member is responding positively to therapy as evidenced by both of the following (a and b): a) at least a 2 mm or greater reduction in proptosis AND b) at least a 2 point or greater reduction in CAS from baseline. Member does not require surgical ophthalmological intervention. Member has not received 8 or more infusions lifetime (including the initial 10 mg/kg first infusion).

### **Age Restrictions:**

#### **Prescriber Restrictions:**

TED: Prescribed by or in consultation with an ophthalmologist.

# **Coverage Duration:**

6 months.

#### Other Criteria:

TED: Failure of a systemic corticosteroid, unless clinically significant adverse effects are experienced or all are contraindicated.

Prior Authorization Group Description:
TETRABENAZINE
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a neurologist.
Coverage Duration:
12 months.
Other Criteria:

### Medicare Part D - 2020

## **Prior Authorization Group Description:**

**TIBSOVO** 

### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

### Off Label Uses:

# **Exclusion Criteria:**

### **Required Medical Information:**

Presence of an isocitrate dehydrogenase-1 (IDH1) mutation. For newly diagnosed acute myeloid leukemia (AML), member is age 60 years or older OR medical justification supports inability to use intensive induction therapy.

#### **Age Restrictions:**

### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

### **Coverage Duration:**

12 months.

# Other Criteria:

For age less than 60 years where medical justification does not support inability to use intensive induction therapy, disease has relapsed or is refractory following treatment with standard antineoplastic induction agents (e.g., cytarabine, idarubicin, daunorubicin, cladribine, midostaurin).

### Medicare Part D - 2020

## **Prior Authorization Group Description:**

**TOLSURA** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

# Off Label Uses:

Hematologic malignancy for prophylaxis of aspergillosis or candidiasis.

# **Exclusion Criteria:**

# **Required Medical Information:**

### **Age Restrictions:**

#### **Prescriber Restrictions:**

# **Coverage Duration:**

HISTOPLASMOSIS: 6 weeks. ASPERGILLOSIS: 3 months. BLASTOMYCOSIS, HEMATOLOGIC MALIGNANCY: 6 months.

# Other Criteria:

ALL INDICATIONS: Failure of generic itraconazole capsule, unless contraindicated or clinically significant adverse effects are experienced. ASPERGILLOSIS: Failure of voriconazole, unless contraindicated or clinically significant adverse effects are experienced. HEMATOLOGIC MALIGNANCY: For candidiasis prophylaxis, failure of fluconazole, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:
TREMFYA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a rheumatologist or dermatologist.
Coverage Duration:
12 months.
Other Criteria:
Failure of ONE of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine, or acitretin.

# Medicare Part D - 2020

TRIHEXYPHENIDYL

# **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

# **Age Restrictions:**

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

# **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

# Other Criteria:

Parkinsons disease/Parkinsonism: Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: amantadine, levodopa/carbidopa, entacapone, pramipexole, ropinirole, selegiline.

#### Medicare Part D - 2020

### **Prior Authorization Group Description:**

**TRIKAFTA** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

### **Required Medical Information:**

CYSTIC FIBROSIS: Diagnosis of cystic fibrosis (CF) confirmed by both of the following (a and b): a) Clinical symptoms consistent with CF in at least one organ system, or positive newborn screen or genetic testing for siblings of patients with CF, AND b) Evidence of cystic fibrosis transmembrane conductance regulator (CFTR) dysfunction confirmed by one of the following (i or ii): i) Evidence of clinical severity as defined by an average sweat chloride greater than 60 mmol/L, OR ii) Genetic testing confirming the presence of two disease-causing mutations in CFTR gene, one from each parent allele, and one of which is a F508del mutation. Confirmation that pulmonary function tests, performed within the last 90 days, show a percent predicted forced expiratory volume in 1 second (ppFEV1) that is between 40-90%. Trikafta is not prescribed concurrently with other CFTR modulators (e.g., Orkambi, Kalydeco, Symdeko). CONTINUATION OF THERAPY, CYSTIC FIBROSIS: For members that received at least 12 weeks of therapy, member is responding positively to therapy as evidenced by stabilization in ppFEV1 if baseline was 70% or greater or increase in ppFEV1 if baseline was less than 70%.

#### **Age Restrictions:**

CYSTIC FIBROSIS: Age greater than or equal to 12 years.

#### **Prescriber Restrictions:**

CYSTIC FIBROSIS: Prescribed by or in consultation with a pulmonologist.

### **Coverage Duration:**

Initial: 4 months. Reauthorization: 12 months.

Prior Authorization Group Description:
TUKYSA
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.
Coverage Duration:
12 months.
Other Criteria:

Prior Authorization Indication: All medically accepted indications not otherwise excluded from Part D.  Diff Label Uses: Exclusion Criteria: Required Medical Information: Age Restrictions: Prescriber Restrictions: Prescribed by or in consultation with an oncologist.  Coverage Duration: 2 months.	<b>Prior Authorization Group Description:</b>	
All medically accepted indications not otherwise excluded from Part D.  Diff Label Uses:  Exclusion Criteria:  Required Medical Information:  Age Restrictions:  Prescriber Restrictions:  Prescribed by or in consultation with an oncologist.  Coverage Duration:  2 months.	TURALIO	
Off Label Uses: Exclusion Criteria:  Required Medical Information:  Age Restrictions: Prescriber Restrictions: Prescribed by or in consultation with an oncologist.  Coverage Duration: 2 months.	Prior Authorization Indication:	
Required Medical Information:  Age Restrictions:  Prescriber Restrictions:  Prescribed by or in consultation with an oncologist.  Coverage Duration:  2 months.	All medically accepted indications not otherwise excluded from Part D.	
Required Medical Information:  Age Restrictions:  Prescriber Restrictions:  Prescribed by or in consultation with an oncologist.  Coverage Duration:  2 months.	Off Label Uses:	
Age Restrictions:  Prescriber Restrictions:  Prescribed by or in consultation with an oncologist.  Coverage Duration:  2 months.	Exclusion Criteria:	
Prescriber Restrictions: Prescribed by or in consultation with an oncologist.  Coverage Duration:  2 months.	Required Medical Information:	
Prescribed by or in consultation with an oncologist.  Coverage Duration:  2 months.	Age Restrictions:	
2 months.	Prescriber Restrictions:	
2 months.	Prescribed by or in consultation with an oncologist.	
	Coverage Duration:	
Other Criteria:	12 months.	
	Other Criteria:	

# Medicare Part D - 2020

# **Prior Authorization Group Description:**

**TYMLOS** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

### Off Label Uses:

# **Exclusion Criteria:**

### **Required Medical Information:**

Total duration of therapy on parathyroid hormone (PTH) analogs (e.g., Tymlos, Forteo) has not exceeded 2 years.

# **Age Restrictions:**

#### **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

#### Other Criteria:

Member meets one of the following (a, b, or c): a) Failure of a bisphosphonate (e.g., alendronate) unless contraindicated or clinically significant adverse effects are experienced. OR b) Bone mineral density T-score at hip or spine is -3.5 or less. OR c) Bone mineral density T-score at hip or spine is -2.5 or less with a history of major osteoporotic fracture of the hip, spine, forearm, wrist, or humerus.

### Medicare Part D - 2020

# **Prior Authorization Group Description:**

**TYSABRI** 

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

# **Exclusion Criteria:**

### **Required Medical Information:**

# **Age Restrictions:**

### **Prescriber Restrictions:**

MULTIPLE SCLEROSIS: Prescribed by or in consultation with a neurologist. CROHN'S DISEASE: Prescribed by or in consultation with a gastroenterologist.

# **Coverage Duration:**

12 months.

# Other Criteria:

RELAPSING-REMITTING MULTIPLE SCLEROSIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Aubagio, Tecfidera, Gilenya, Avonex, Betaseron, Plegridy, glatiramer, Copaxone, Glatopa, Extavia or Rebif. CROHN'S DISEASE: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Humira or Remicade.

<b>Prior Authorization Group Description:</b>
UPTRAVI
<b>Prior Authorization Indication:</b>
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

Prior Authorization Group Description:
VALCHLOR
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of one of the following skin-directed therapies unless contraindicated or clinically significant adverse effe are experienced: topical corticosteroid (e.g., betamethasone, clobetasol), topical retinoid (e.g., Targretin, Avage, Fabior, Tazorac), topical imiquimod (Aldara).

# Medicare Part D - 2020

VALTOCO

# **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

Diagnosis of partial or generalized epilepsy.

# **Age Restrictions:**

6 years of age or older.

# **Prescriber Restrictions:**

Prescribed by or in consultation with a neurologist.

# **Coverage Duration:**

12 months.

# Other Criteria:

Medical justification supports inability to use diazepam rectal gel (e.g., contraindications to excipients).

# **Medicare Part D – 2020**

Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
C. Diff diarrhea: 14 days. Staph enterocolitis: 10 days. Recurrent C. Diff: 12 weeks.
Other Criteria:

**Prior Authorization Group Description:** 

VANCOCIN

### Medicare Part D - 2020

## **Prior Authorization Group Description:**

**VENCLEXTA** 

### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

### Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

AML: Age 60 years or greater, OR medical justification supports inability to use intensive induction chemotherapy. Prescribed in combination with azacitidine, decitabine, or low-dose cytarabine.

#### **Age Restrictions:**

### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

### **Coverage Duration:**

12 months.

# Other Criteria:

MANTLE CELL LYMPHOMA: Failure of at least one previous therapy (e.g., a Rituxan based regimen), unless contraindicated or clinically significant adverse effects are experienced. CLL/SLL: Request meets one of the following (a or b): a) Prescribed in combination with Gazyva as first-line therapy OR b) Failure of at least one previous therapy (e.g., Imbruvica, Campath, or high-dose methylprednisolone with Rituxan), unless contraindicated or clinically significant adverse effects are experienced.

<b>Prior Authorization Group Description:</b>
VERSACLOZ
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Psychotic disorder associated with Parkinson's disease.
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of clozapine (Clozaril) or FazaClo, unless contraindicated or clinically significant adverse effects are experienced.

### Medicare Part D - 2020

# **Prior Authorization Group Description:**

**VERZENIO** 

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

### Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

Hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced or metastatic breast cancer.

#### **Age Restrictions:**

### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

### **Coverage Duration:**

12 months.

# Other Criteria:

Prescribed as a single agent or in combination with an aromatase inhibitor (e.g., letrozole, anastrozole or exemestane) or fulvestrant. For men receiving an aromatase inhibitor: Prescribed in combination with an agent that suppresses testicular steroidogenesis (e.g., gonadotropin-releasing hormone agonists).

# Medicare Part D – 2020

<b>Prior Authorization Group Description:</b>
VINBLASTINE
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Confirmation that vinblastine is being used as palliative therapy.
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with an oncologist or hematologist.

**Coverage Duration:** 

12 months.

# Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>	

VINCRISTINE

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

Patients with the demyelinating form of Charcot-Marie-Tooth syndrome.

# **Required Medical Information:**

# **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

# **Coverage Duration:**

12 months.

# Medicare Part D - 2020

VITRAKVI

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

Known acquired tropomyosin receptor kinase resistance mutation.

# **Required Medical Information:**

Confirmation of positive neurotrophic receptor tyrosine kinase gene fusion mutation.

# **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# Medicare Part D - 2020

<b>Prior Authorization</b>	<b>Group Description:</b>	

**VIZIMPRO** 

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

Disease is positive for a sensitizing EGFR mutation (e.g., exon 19 deletion or insertion, exon 21 point mutation (L858R, L861Q), exon 18 point mutation (G719X), exon 20 point mutation (S768I)).

# **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

### Medicare Part D - 2020

# **Prior Authorization Group Description:**

VOSEVI

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

### Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

If cirrhosis is present, confirmation of Child-Pugh A status. Refer to the document, "Recommendations for Testing, Managing, and Treating Hepatitis C," by AASLD-IDSA available at http://www.hcvguidelines.org for drug regimen and duration of treatment based on genotype, treatment status, previous drug regimens used, past medical history and comorbidities.

# **Age Restrictions:**

### **Prescriber Restrictions:**

Prescribed by or in consultation with a gastroenterologist, hepatologist or infectious disease physician.

### **Coverage Duration:**

12 weeks.

# Other Criteria:

Criteria will be applied consistent with current AASLD-IDSA guidance.

# Medicare Part D – 2020

**Prior Authorization Group Description:** 

VOTRIENT
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with an oncologist.
Coverage Duration:
12 months.
Other Criteria:

Prior Authorization Group Description:
VRAYLAR
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:
Failure of TWO of the following atypical antipsychotics, unless contraindicated or clinically significant adverse effects are experienced: aripiprazole, ziprasidone, quetiapine, olanzapine, risperidone.

# Medicare Part D – 2020

VUMERITY
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
MULTIPLE SCLEROSIS: Prescribed by or in consultation with a neurologist.
Coverage Duration:
12 months.
Other Criteria:

**Prior Authorization Group Description:** 

# Medicare Part D - 2020

# **Prior Authorization Group Description:**

VYONDYS 53

### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

DUCHENNE MUSCULAR DYSTROPHY (DMD): DMD with mutation amenable to exon 53 skipping confirmed by genetic testing.

### **Age Restrictions:**

# **Prescriber Restrictions:**

DMD: Prescribed by or in consultation with a neurologist.

# **Coverage Duration:**

6 months.

# Other Criteria:

DMD: Currently stable on an oral corticosteroid regimen (e.g., prednisone), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:
WAKIX
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
NARCOLEPSY: Prescribed by or in consultation with a neurologist.
Coverage Duration:
12 months.
Other Criteria:
NARCOLEPSY: Failure of Sunosi, unless contraindicated or clinically significant adverse effects are experienced.

# Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>		

XALKORI

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

NON-SMALL CELL LUNG CANCER: Disease is ALK, ROS1, or MET positive. INFLAMMATORY MYOFIBROBLASTIC TUMOR, ANAPLASTIC LARGE CELL LYMPHOMA: Disease is ALK-positive.

# **Age Restrictions:**

# **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

# Medicare Part D - 2020

# **Prior Authorization Group Description:**

**XATMEP** 

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

# **Age Restrictions:**

Less than 18 years of age.

# **Prescriber Restrictions:**

ACUTE LYMPHOCYTIC LEUKEMIA: Prescribed by or in consultation with an oncologist or hematologist. POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist.

# **Coverage Duration:**

12 months.

### Other Criteria:

Medical justification as to why member cannot use methotrexate tablets.

Prior Authorization Group Description:
XCOPRI
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a neurologist.
Coverage Duration:
12 months.
Other Criteria:
Failure of two of the following generic antiepileptic drugs, unless contraindicated or clinically significant adverse effects are experienced: lamotrigine, topiramate, oxcarbazepine, carbamazepine, phenytoin, valproic acid, divalproex sodium, felbamate, gabapentin, levetiracetam, pregabalin, tiagabine, zonisamide.

#### Medicare Part D - 2020

# Prior Authorization Group Description:

**XELJANZ** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

**Off Label Uses:** 

**Exclusion Criteria:** 

#### **Required Medical Information:**

**Age Restrictions:** 

#### **Prescriber Restrictions:**

RHEUMATOID ARTHRITIS: Prescribed by or in consultation with a rheumatologist. PSORIATIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist or dermatologist. ULCERATIVE COLITIS (IMMEDIATE-RELEASE ONLY): Prescribed by or in consultation with a gastroenterologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. PSORIATIC ARTHRITIS: Failure of one TNF inhibitor (e.g., Humira, Enbrel, Cimzia, Simponi, Simponi Aria, Remicade, Inflectra), unless predominantly axial disease, contraindicated, or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

Prior Authorization	Group	<b>Description:</b>
---------------------	-------	---------------------

**XENICAL** 

#### **Prior Authorization Indication:**

For obesity management including weight loss and weight maintenance when used in conjunction with a reduced-calorie diet or to reduce the risk for weight regain after prior weight loss in obese patients with an initial body mass index (BMI) of 30 kg/m<sup>2</sup> or greater (obese) or 27 kg/m<sup>2</sup> or greater (overweight) in the presence of at least one weight-related comorbidity such as hypertension, type 2 diabetes mellitus, or dyslipidemia.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

BMI is greater than or equal to 30 kg/m<sup>2</sup> OR BMI is greater than or equal to 27 kg/m<sup>2</sup> with one or more of the following severe co-morbid conditions 1. Coronary artery/heart disease 2. Diabetes 3. Dyslipidemia 4. Hypertension 5. Obstructive sleep apnea. CONTINUATION OF THERAPY: Confirmation of a 5-10 pound weight loss during the first 6 month treatment period. Therapy beyond the first year can be authorized every 6 months with confirmation of weight maintenance.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

#### **Coverage Duration:**

6 months.

#### Medicare Part D - 2020

<b>Prior Authorization</b>	Group	<b>Description:</b>
----------------------------	-------	---------------------

**XENLETA** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

Request is for continuation of therapy initiated in an acute care hospital from which member was discharged OR culture and sensitivity (C&S) report for the current infection shows isolated pathogen is an organism susceptible to Xenleta, unless provider confirms that obtaining a C&S report is not feasible.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

#### **Coverage Duration:**

7 days.

#### Other Criteria:

For members initiating Xenleta therapy outside of an acute care hospital, one of the following (a, b, or c): a) If a C&S report is available: Failure of 2 antibiotics to which the isolated pathogen is susceptible (if available) per C&S report, unless all are contraindicated or clinically significant adverse effects are experienced. b) C&S report shows resistance or lack of susceptibility of the isolated pathogen to all formulary antibiotics FDA-approved for member's diagnosis. c) If provider confirms that obtaining a C&S report is not feasible: Failure of 2 antibiotics indicated for member's diagnosis (if available), unless all are contraindicated or clinically significant adverse effects are experienced.

# Medicare Part D – 2020

<b>Prior Authorization Group Description:</b>
XEOMIN
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D - 2020

Prior	Authorization	Groun	Descri	ntion:	
	I LUCIIOI ILUCIOII	Group	Descri	J 41 0 11 4	

**XERMELO** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

# **Required Medical Information:**

Prescribed in combination with a somatostatin analog (e.g., octreotide, lanreotide) unless contraindicated or clinically significant adverse effects are experienced.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

# **Coverage Duration:**

12 months.

# Other Criteria:

Failure of a somatostatin analog (e.g., octreotide, lanreotide) unless contraindicated or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**XOLAIR** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

ASTHMA: Positive skin test or in vitro reactivity to a perennial aeroallergen AND immunoglobulin E (IgE) level greater than or equal to 30 IU/mL.

#### **Age Restrictions:**

ASTHMA: 6 years of age or older. CHRONIC IDIOPATHIC URTICARIA: 12 years of age or older.

#### **Prescriber Restrictions:**

ASTHMA: Prescribed by or in consultation with a pulmonologist, immunologist, or allergist. CHRONIC IDIOPATHIC URTICARIA: Prescribed by or in consultation with an allergist, dermatologist, or immunologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

ASTHMA: Failure of one inhaled corticosteroid (e.g., beclomethasone, budesonide, flunisolide, fluticasone, mometasone, ciclesonide), unless contraindicated or clinically significant adverse effects are experienced. CHRONIC IDIOPATHIC URTICARIA: Failure of one H1 Antihistamine (e.g., levocetirizine or desloratadine), unless contraindicated or clinically significant adverse effects are experienced.

# Medicare Part D - 2020

Prior Authorization	Group Description:	
TIOI TIMENOTIZMENT	Group Description.	

**XOSPATA** 

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

Confirmation of the presence of a FLT3 mutation.

# **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

# **Coverage Duration:**

12 months.

#### Medicare Part D - 2020

Prior	Authorization	Group	<b>Description:</b>

**XPOVIO** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

#### **Exclusion Criteria:**

# **Required Medical Information:**

MULTIPLE MYELOMA: Member has received at least 4 prior lines of therapy that include all of the following (a, b, and c): a) Two proteasome inhibitors (e.g., bortezomib, Kyprolis, Ninlaro), b) Two immunomodulatory agents (e.g., Revlimid, pomalidomide, Thalomid), c) One anti-CD38 monoclonal antibody (e.g., Darzalex).

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or hematologist.

#### **Coverage Duration:**

12 months.

# Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>		
XTANDI		

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

# Off Label Uses:

# **Exclusion Criteria:**

# **Required Medical Information:**

PROSTATE CANCER: Concurrent use of a gonadotropin-releasing hormone (GnRH) analog or past bilateral orchiectomy.

# **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or urologist.

# **Coverage Duration:**

12 months.

# Medicare Part D - 2020

#### **Prior Authorization Group Description:**

YERVOY

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

HEPATOCELLULAR CARCINOMA (new starts only): Member has not had previous treatment with a checkpoint inhibitor (e.g., Opdivo, Keytruda, Tecentriq, Imfinzi). NON-SMALL CELL LUNG CANCER (new starts only): Member has not previously progressed on a PD-1/PD-L1 inhibitor (e.g., Opdivo, Keytruda, Tecentriq, Imfinzi).

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

SMALL CELL LUNG CANCER, MALIGNANT PLEURAL MESOTHELIOMA: Failure of a platinum-containing regimen (e.g., cisplatin, carboplatin), unless contraindicated or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

# Prior Authorization Group Description: YONSA Prior Authorization Indication: All medically accepted indications not otherwise excluded from Part D. Off Label Uses: Exclusion Criteria: Required Medical Information: Age Restrictions: Prescriber Restrictions: Prescribed by or in consultation with an oncologist or urologist. Coverage Duration: 12 months. Other Criteria: Medical justification supports inability to use Zytiga. Prescribed in combination with methylprednisolone. Member

has previously had bilateral orchiectomy, failed androgen deprivation therapy (ADT) or will use ADT concurrently

with Yonsa.

#### Medicare Part D - 2020

# Prior Authorization Group Description:

**ZALTRAP** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

**Exclusion Criteria:** 

#### **Required Medical Information:**

**Age Restrictions:** 

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

# **Coverage Duration:**

12 months.

#### Other Criteria:

Prescribed in combination with irinotecan or FOLFIRI (5-fluorouracil, leucovorin, and irinotecan). Previous treatment with one of the following: oxaliplatin-containing regimen (e.g., FOLFIRI, FOLFOX [leucovorin, 5-fluorouracil, oxaliplatin], CapeOX [capecitabine, oxaliplatin]) OR 5-fluorouracil and leucovorin containing regimen OR capecitabine containing regimen.

# Medicare Part D – 2020

Prior Authorization Group Description:
ZARXIO
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Myelodysplastic syndrome. Neutropenia in patients with HIV/AIDS. Hematopoietic syndrome of acute radiation syndrome.
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
12 months.
Other Criteria:

# Medicare Part D – 2020

Prior Authorization Group Description:
ZEJULA
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
OVARIAN, FALLOPIAN, OR PRIMARY PERITONEAL CANCER: Prescribed by or in consultation with an oncologist.
Coverage Duration:
12 months.
Other Criteria:

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**ZELBORAF** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

MELANOMA, ERDHEIM-CHESTER DISEASE: Positive for the BRAF V600 mutation. NON-SMALL CELL LUNG CANCER, COLORECTAL CANCER: Positive for the BRAF V600E mutation. DIFFERENTIATED THYROID CARCINOMA: Positive for the BRAF mutation.

#### **Age Restrictions:**

# **Prescriber Restrictions:**

ERDHEIM-CHESTER DISEASE, HAIRY CELL LEUKEMIA: Prescribed by or in consultation with an oncologist or hematologist. ALL OTHER ONCOLOGY INDICATION: Prescribed by or in consultation with an oncologist.

#### **Coverage Duration:**

12 months.

#### Other Criteria:

NON-SMALL CELL LUNG CANCER: Failure of Tafinlar or Mekinist, unless contraindicated or clinically significant adverse effects are experienced. COLORECTAL CANCER: Failure of irinotecan or platinum-based therapy (e.g., oxaliplatin), unless contraindicated or clinically significant adverse effects are experienced.

#### Medicare Part D - 2020

#### **Prior Authorization Group Description:**

**ZEPATIER** 

#### **Prior Authorization Indication:**

All FDA-approved indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

#### **Required Medical Information:**

If cirrhosis is present, confirmation of Child-Pugh A status. For genotype 1a, confirmation of presence or absence of NS5A resistance-associated polymorphisms. Refer to the document, "Recommendations for Testing, Managing, and Treating Hepatitis C," by AASLD-IDSA available at http://www.hcvguidelines.org for drug regimen and duration of treatment based on genotype, treatment status, previous drug regimens used, past medical history and comorbidities.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with a gastroenterologist, hepatologist or infectious disease physician.

#### **Coverage Duration:**

12 to 16 wks based on genotype, presence of NS5A resistance-associated polymorphisms, prior treatment.

# Medicare Part D – 2020

**Prior Authorization Group Description:** 

CDI recurrences, including a pulsed vancomycin regimen.

ZINPLAVA
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Confirmation of positive Clostridium difficile test.
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
4 weeks.
Other Criteria:
Will receive or is currently receiving antibacterial drug treatment for Clostridium difficile infection (e.g., metronidazole, vancomycin, fidaxomicin) concomitantly with Zinplava. Has received appropriate treatment for past

# Medicare Part D – 2020

Prior Authorization Group Description:
ZULRESSO
Prior Authorization Indication:
All FDA-approved indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
No more than 6 months have passed since member has given birth.
Age Restrictions:
Prescriber Restrictions:
Coverage Duration:
4 weeks.
Other Criteria:
Failure of one of the following oral antidepressants, unless contraindicated or clinically significant adverse effects are experienced: selective serotonin reuptake inhibitor (SSRI), serotonin-norepinephrine reuptake inhibitor (SNRI) tricyclic antidepressant (TCA), bupropion, mirtazapine.

# **Medicare Part D – 2020**

ZYDELIG
Prior Authorization Indication:
All medically accepted indications not otherwise excluded from Part D.
Off Label Uses:
Exclusion Criteria:
Required Medical Information:
Age Restrictions:
Prescriber Restrictions:
Prescribed by or in consultation with a hematologist or oncologist.
<b>Coverage Duration:</b>
12 months.
Other Criteria:

**Prior Authorization Group Description:** 

# Medicare Part D - 2020

**ZYKADIA** 

#### **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

#### Off Label Uses:

#### **Exclusion Criteria:**

# **Required Medical Information:**

NON-SMALL CELL LUNG CANCER: Disease is ALK or ROS1 positive. If disease is ROS1 positive, Zykadia is prescribed as first-line therapy. INFLAMMATORY MYOFIBROBLASTIC TUMOR: Disease is ALK-positive.

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist.

#### **Coverage Duration:**

12 months.

# Medicare Part D - 2020

<b>Prior Authorization Group Description:</b>		

# **Prior Authorization Indication:**

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

**ZYTIGA** 

**Exclusion Criteria:** 

# **Required Medical Information:**

#### **Age Restrictions:**

#### **Prescriber Restrictions:**

Prescribed by or in consultation with an oncologist or urologist.

# **Coverage Duration:**

12 months.

#### Other Criteria:

Prescribed in combination with prednisone. Member has previously had bilateral orchiectomy, failed androgen deprivation therapy (ADT) or will use ADT concurrently with abiraterone.

ACTEMRA SC1	CINQAIR4	18
ACTIQ2	CLADRIBINE4	19
ACYCLOVIR	CLOMIPRAMINE5	50
ADAKVEO 4	COMETRIQ5	51
ADCIRCA 5	COPIKTRA 5	52
ADEMPAS6	COTELLIC5	53
AFINITOR7	CRYSVITA5	54
AIMOVIG8	CYCLOBENZAPRINE HCL5	55
AJOVY9	CYTARABINE	56
ALECENSA	DAURISMO5	57
ALLI11	DICLOFENAC GEL	58
ALUNBRIG12	DIETHYLPROPION5	59
AMITRIPTYLINE	DIPYRIDAMOLE6	50
AMITRIPTYLINE/CHLORDIAZEPOXIDE 14	DISOPYRAMIDE6	
AMITRIPTYLINE/PERPHENAZINE	DOPTELET6	52
AMPHOTERICIN B	DOXEPIN	
AMPYRA	DOXEPIN CREAM	54
ANTIHISTAMINES	ELIDEL	
ANTIHISTAMINE COMBINATIONS 19	EMEND 40 MG	
ARANESP	EMFLAZA6	
ARIKAYCE21	EMGALITY	
AUBAGIO	ENBREL	
AYVAKIT23	ENDARI	
BALVERSA	ENTYVIO	
BAXDELA	EPCLUSA	
BELEODAQ	EPIDIOLEX7	
BELSOMRA	EPOETIN	
BENLYSTA	ERGOLOID MESYLATES	75
BENZTROPINE	ERLEADA	76
BEOVU30	ESBRIET	77
BLEOMYCIN	ESTROGENS(Divigel, Climara, Mimvey,	
BOSULIF32	Prempro, Amabelz, Estrace, Activella, Femhro Mimvey Lo, Lopreeza, Premarin, Premphase	t,
BOTOX33	)	78
BRAFTOVI34	EXONDYS 51	
BRIVIACT	FARYDAK8	
BRUKINSA	FASENRA8	
C1 ESTERASE INHIBITOR	FERRIPROX8	
CABLIVI38	FINTEPLA8	
CABOMETYX	FIORINAL WITH CODEINE	
CALQUENCE	FIRAZYR8	
CAPLYTA41	FIRDAPSE8	
CAPRELSA42	FLECTOR8	
CAYSTON	FLUOROURACIL8	
CERDELGA	FORTEO	
CEREZYME	GALAFOLD9	
CHLORZOXAZONE	GANCICLOVIR9	
CHORIONIC GONADOTROPIN 47	GATTEX9	<del>)</del> 2

GAVRETO93	LENVIMA140
GILENYA	LEUKINE141
GILOTRIF95	LIDODERM142
GLATIRAMER96	LONSURF143
GLIMEPIRIDE	LORBRENA144
GLYBURIDE98	LOTRONEX
GLYBURIDE/METFORMIN	LUCEMYRA146
GRANIX	LYNPARZA TABLET147
HARVONI101	MAVENCLAD148
HERCEPTIN102	MAVYRET149
HETLIOZ103	MAYZENT150
HUMAN GROWTH HORMONE104	MEGACE151
HUMIRA105	MEGACE ES
HYDROCODONE106	MEKINIST
HYDROXYZINE HCL INJECTION	MEKTOVI
HYDROXYZINE HCL ORAL	METAXALONE
HYDROXYZINE PAMOATE109	METHAMPHETAMINE
ICLUSIG	METHOCARBAMOL
IDHIFA	METHOTREXATE INJ
ILARIS112	MIRVASO
ILUMYA113	MOZOBIL
IMATINIB114	MULPLETA161
IMBRUVICA115	NAMENDA
IMIPRAMINE116	NATPARA163
INDOMETHACIN117	NAYZILAM
INFLECTRA118	NERLYNX
INGREZZA119	NEULASTA166
INLYTA120	NEUPOGEN
INQOVI	NINLARO
INREBIC	NIVESTYM
INTERFERON BETA-1A123	NORTHERA170
INTERFERON BETA-1B	NUBEQA171
INTUNIV	NUCALA
JAKAFI	NUEDEXTA
JUXTAPID127	NUPLAZID174
JYNARQUE128	NUZYRA175
KADCYLA	OCALIVA
KALYDECO130	OCREVUS
KETOROLAC TROMETHAMINE	ODOMZO178
KEVZARA	OFEV
KISQALI(Kisqali Femara Co-Pack, Kisqali)133	OLUMIANT
KORLYM	ONUREG
KOSELUGO	OPSUMIT
KUVAN	ORENITRAM
LATUDA137	ORILISSA
LAZANDA	ORKAMBI185
LEMTRADA139	OXBRYTA186

OXERVATE	STELARA IV	234
PALYNZIQ188	STELARA SC	235
PEMAZYRE	STIVARGA	236
PENNSAID	STRENSIQ	237
PERSERIS	SUBSYS	238
PHENOBARBITAL	SUNOSI	239
PHENTERMINE	SURMONTIL	240
PIQRAY	SYMDEKO	241
PRALUENT195	SYMLINPEN	242
PRETOMANID	SYMPAZAN	243
PREVYMIS197	TABRECTA	244
PROLASTIN C	TAFAMIDIS	245
PROLIA199	TAGRISSO	246
PROMACTA200	TAKHZYRO	247
PROTOPIC	TALZENNA	248
PROVIGIL202	TARCEVA	249
PURIXAN	TARGRETIN GEL	250
QINLOCK	TASIGNA	251
QSYMIA	TAVALISSE	252
QUALAQUIN	TAZVERIK	253
RADICAVA	TECENTRIQ	254
RAYALDEE	TECFIDERA	255
REBLOZYL	TEGSEDI	256
REMICADE	TENEX	257
RENFLEXIS	TEPEZZA	258
REPATHA212	TETRABENAZINE	259
RETEVMO213	TIBSOVO	260
REVATIO	TOLSURA	261
REVCOVI	TREMFYA	262
REVLIMID	TRIHEXYPHENIDYL	263
REXULTI217	TRIKAFTA	264
RINVOQ	TUKYSA	265
ROZLYTREK219	TURALIO	266
RUBRACA	TYMLOS	267
RUZURGI	TYSABRI	268
RYDAPT222	UPTRAVI	269
SECUADO	VALCHLOR	270
SILIQ224	VALTOCO	271
SIMPONI(prefilled syringe, auto-injector)225	VANCOCIN	272
SIMPONI ARIA	VENCLEXTA	273
SKYRIZI227	VERSACLOZ	274
SOMA	VERZENIO	275
SOMAVERT	VINBLASTINE	276
SOVALDI	VINCRISTINE	277
SPRAVATO231	VITRAKVI	278
SPRITAM	VIZIMPRO	279
SPRYCEL	VOSEVI	280

VOTRIENT	281
VRAYLAR	282
VUMERITY	283
VYONDYS 53	284
WAKIX	285
XALKORI	286
XATMEP	287
XCOPRI	288
XELJANZ	. 289
XENICAL	. 290
XENLETA	291
XEOMIN	292
XERMELO	293
XOLAIR	294
XOSPATA	295
XPOVIO	296
XTANDI	297
YERVOY	298
YONSA	299
ZALTRAP	300
ZARXIO	301
ZEJULA	302
ZELBORAF	303
ZEPATIER	304
ZINPLAVA	305
ZULRESSO	306
ZYDELIG	307
ZYKADIA	308
ZYTIGA	309