Prior Authorization Protocol

Medicare Part D - 2020

Prior Authorization Group Description:

ABSTRAL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Age 18 or greater.

Prescriber Restrictions:

Coverage Duration:

Through the end of the Plan contract year.

Other Criteria:

Patient is already taking and is tolerant to around-the-clock opioid therapy. Patients are considered opioid tolerant when taking another opioid daily for a week or longer (for example, at least 60 mg of oral morphine per day or an equianalgesic dose of another opioid).

Prior Authorization Group Description:

ACTEMRA IV

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

RHEUMATOID ARTHRITIS: Prescribed by or in consultation with a rheumatologist. JUVENILE IDIOPATHIC ARTHRITIS: Prescribed by or in consultation with a dermatologist, rheumatologist, or gastroenterologist.

Coverage Duration:

Rheumatoid Arthritis, Juvenile Idiopathic Arthritis: 12 months. Cytokine Release Syndrome: 3 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of Remicade and one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin.

Prior Authorization Group Description:

ACTEMRA SC

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

RHEUMATOID ARTHRITIS, GIANT CELL ARTERITIS: Prescribed by or in consultation with a rheumatologist. JUVENILE IDIOPATHIC ARTHRITIS: Prescribed by or in consultation with a dermatologist, rheumatologist, or gastroenterologist.

Coverage Duration:

12 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. GIANT CELL ARTERITIS: Failure of methotrexate or azathioprine, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

ACTHAR HP

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist, nephrologist, rheumatologist, dermatologist, ophthalmologist, or allergist/immunologist as appropriate.

Coverage Duration:

MS,RA,JUVENILE RA,PSORIATIC ARTHRITIS,ANKYLOSING SPONDYLITIS:1 month.All other indications:3 months.

Other Criteria:

MULTIPLE SCLEROSIS: Member is being treated with a relapsing remitting multiple sclerosis agent (e.g., Avonex, Betaseron, Copaxone, Gilenya) AND Failure of corticosteroid therapy for acute exacerbations of multiple sclerosis, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

ACTIQ

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

16 years and older.

Prescriber Restrictions:

Coverage Duration:

Through the end of the Plan contract year.

Other Criteria:

Member is already taking and is tolerant to around-the-clock opioid therapy. Members are considered opioid tolerant when taking another opioid daily for a week or longer (for example, at least 60 mg of oral morphine per day or an equianalgesic dose of another opioid).

Prior Authorization Group Description:

ACYCLOVIR

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

ADAKVEO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

SICKLE CELL DISEASE: Disease is associated with one of the following genotypes: Homozygous hemoglobin S, Hemoglobin S beta 0-thalassemia, Hemoglobin S beta+ thalassemia, Hemoglobin SC. Member has a hemoglobin level of at least 4 g/dL. Member meets one of the following (a or b): a) Member experienced at least 2 vaso-occlusive crises (VOC) within the past 6 months while on hydroxyurea, OR b) Member has intolerance or contraindication to hydroxyurea and has experienced at least 2 VOC within the past 12 months. Confirmation of baseline incidence of VOC over the last 12 months. Adakveo is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced. Adakveo is not prescribed concurrently with Oxbryta. CONTINUATION OF THERAPY, SICKLE CELL DISEASE: Member is responding positively to therapy as evidenced by an improvement in the incidence of VOC from baseline. Adakveo is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse of VOC from baseline. Adakveo is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse of VOC from baseline. Adakveo is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced. Adakveo is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced. Adakveo is not prescribed concurrently with Oxbryta.

Age Restrictions:

SICKLE CELL DISEASE: Age greater than or equal to 16 years.

Prescriber Restrictions:

SICKLE CELL DISEASE: Prescribed by or in consultation with a hematologist.

Coverage Duration:

6 months.

Other Criteria:

SICKLE CELL DISEASE: Failure of L-glutamine, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

ADCIRCA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Members on concomitant nitrates (e.g., Nitrodur, Nitrobid, Nitrostat, Isordil, Ismo). Members on concomitant guanylate cyclase stimulator, such as riociguat (Adempas).

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

ADEMPAS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Members on concomitant phosphodiesterase (PDE) inhibitors (e.g., sildenafil, tadalafil, vardenafil, dipyridamole or theophylline) or nitrates (e.g., Nitrodur, Nitrobid, Nitrostat, Isordil, Ismo).

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

AFINITOR

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

BREAST CANCER: hormone-receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative recurrent or metastatic disease. TUBEROUS SCLEROSIS COMPLEX WITH SUBEPENDYMAL GIANT CELL ASTROCYTOMA: Member is not a candidate for curative surgical resection.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist. TUBEROUS SCLEROSIS COMPLEX ASSOCIATED PARTIAL ONSET SEIZURES: Prescribed by or in consultation with an oncologist or neurologist.

Coverage Duration:

12 months.

Other Criteria:

RENAL CELL CARCINOMA WITH CLEAR CELL HISTOLOGY: Failure of one prior therapy (e.g., Votrient, Sutent), unless contraindicated or clinically significant adverse effects are experienced. BREAST CANCER: Prescribed in combination with exemestane, fulvestrant or tamoxifen AND history of prior endocrine therapy (e.g., letrozole, anastrozole) unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

AIMOVIG

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

One of the following (a or b): a) Member experiences 4 or more migraine days per month for at least 3 months, or b) Diagnosis of chronic migraine and Aimovig is prescribed for prophylaxis. CONTINUATION OF THERAPY: Member has experienced and maintained positive response to therapy as evidenced by a reduction in migraine days per month from baseline.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist, headache, or pain specialist.

Coverage Duration:

Initial: 3 months. Reauthorizations: 6 months.

Other Criteria:

Failure of two of the following oral migraine preventative therapies, each from different therapeutic classes, unless contraindicated or clinically significant adverse effects are experienced: antiepileptic drugs (e.g., divalproex sodium, sodium valproate, topiramate), beta-blockers (e.g., metoprolol, propranolol, timolol), antidepressants (e.g., amitriptyline, venlafaxine).

Prior Authorization Group Description:

AJOVY

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

One of the following (a or b): a) Member experiences 4 or more migraine days per month for at least 3 months, or b) Diagnosis of chronic migraine and Ajovy is prescribed for prophylaxis. CONTINUATION OF THERAPY: Member has experienced and maintained positive response to therapy as evidenced by a reduction in migraine days per month from baseline.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist, headache, or pain specialist.

Coverage Duration:

Initial: 3 months. Reauthorizations: 6 months.

Other Criteria:

Failure of two of the following oral migraine preventative therapies, each from different therapeutic classes, unless contraindicated or clinically significant adverse effects are experienced: antiepileptic drugs (e.g., divalproex sodium, sodium valproate, topiramate), beta-blockers (e.g., metoprolol, propranolol, timolol), antidepressants (e.g., amitriptyline, venlafaxine).

Prior Authorization Group Description:

ALECENSA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Disease is anaplastic lymphoma kinase (ALK) positive.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

ALUNBRIG

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Disease is anaplastic lymphoma kinase (ALK) positive.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

AMANTADINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Member meets one of the following (a or b): a) Medical justification supports inability to use immediate-release amantadine OR b) Failure of immediate-release amantadine unless contraindicated or clinically significant adverse effects are experienced.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

AMITRIPTYLINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Depression: Failure of one of the following generic antidepressants, unless contraindicated or clinically significant adverse effects are experienced: bupropion, bupropion SR, bupropion XL, citalopram, desvenlafaxine succinate, duloxetine, escitalopram, fluoxetine, mirtazapine, sertraline, venlafaxine, or venlafaxine XR.

Prior Authorization Group Description:

AMITRIPTYLINE/CHLORDIAZEPOXIDE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: duloxetine, escitalopram, or venlafaxine XR.

Prior Authorization Group Description:

AMITRIPTYLINE/PERPHENAZINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Depression: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: duloxetine, escitalopram, or venlafaxine XR.

Prior Authorization Group Description:

AMPHOTERICIN B

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Abelcet only: Failure of conventional amphotericin B therapy unless contraindicated or clinically significant adverse effects are experienced. Ambisome when treating patients with Aspergillus species, Candida species and/or Cryptococcus species infections: Failure of conventional amphotericin B therapy unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

AMPYRA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

ANTIHISTAMINES

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Allergic rhinitis: Failure to two of the following, unless contraindicated or clinically significant adverse effects are experienced: levocetirizine, desloratadine, fluticasone propionate nasal suspension, flunisolide nasal solution or triamcinolone acetonide nasal inhaler.

Prior Authorization Group Description:

ANTIHISTAMINE COMBINATIONS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Allergic rhinitis: Failure to two of the following, unless contraindicated or clinically significant adverse effects are experienced: levocetirizine, desloratadine, fluticasone propionate nasal suspension, flunisolide nasal solution or triamcinolone acetonide nasal inhaler.

Prior Authorization Group Description:

ARANESP

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Anemia due to myelodysplastic syndrome. Myelofibrosis-associated anemia.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of Procrit, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

ARIKAYCE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Positive sputum culture after at least 6 consecutive months of a multidrug background regimen therapy (e.g., clarithromycin or azithromycin, ethambutol, and a rifamycin). CONTINUATION OF THERAPY: Confirmation of at least 3 consecutive negative monthly sputum cultures in the first 6 months of therapy or at least 2 consecutive negative monthly sputum cultures in the last 2 months of therapy. Member has not received Arikayce treatment for more than 12 months after converting to negative sputum status.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an infectious disease specialist or pulmonologist.

Coverage Duration:

Initial: 6 months. Reauthorizations: 12 months.

Prior Authorization Group Description:

AUBAGIO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

AUSTEDO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

TARDIVE DYSKINESIA: Development of tardive dyskinesia is secondary to a centrally acting dopamine receptor blocking agent (neuroleptic) (e.g., first- or second-generation antipsychotics such as chlorpromazine or aripiprazole, antiemetics such as promethazine or metoclopramide, the tri-cyclic antidepressant amoxapine).

Age Restrictions:

Prescriber Restrictions:

HUNTINGTON'S DISEASE: Prescribed by or in consultation with a neurologist. TARDIVE DYSKINESIA: Prescribed by or in consultation with a psychiatrist or neurologist.

Coverage Duration:

12 months.

Other Criteria:

HUNTINGTON'S DISEASE: Failure of tetrabenazine, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

AVASTIN

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Age 18 or older.

Prescriber Restrictions:

All cancer indications: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

FOR ALL NON-OPHTHALMOLOGY INDICATIONS: Medical justification supports inability to use Mvasi or Zirabev (e.g., contraindications to the excipients). Non-squamous non-small cell lung cancer: Prescribed in combination with carboplatin and paclitaxel for first line treatment of unresectable, locally advanced, recurrent or metastatic disease. Metastatic renal cell carcinoma: Prescribed in combination with interferon alfa.

Prior Authorization Group Description:

AYVAKIT

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

GASTROINTESTINAL STROMAL TUMOR: Confirmation of PDGFRA exon 18 mutation.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

GASTROINTESTINAL STROMAL TUMOR: If disease is positive for a PDGFRA exon 18 mutation other than D842V, failure of imatinib, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

BALVERSA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Presence of susceptible fibroblast growth factor receptor (FGFR) 3 or FGFR2 genetic alterations.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

Disease has progressed during or following at least one line of platinum-containing chemotherapy.

Prior Authorization Group Description:

BAXDELA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

ACUTE BACTERIAL SKIN AND SKIN STRUCTURE INFECTION (ABSSSI), COMMUNITY-ACQUIRED BACTERIAL PNEUMONIA (CABP): Request is for continuation of therapy initiated in an acute care hospital from which member was discharged OR Both of the following (1 and 2): 1) Current culture and sensitivity (C&S) report shows isolated pathogen is susceptible to delafloxacin, unless provider confirms that obtaining a C&S report is not feasible AND 2) Failure of one fluoroquinolone, unless all are contraindicated or clinically significant adverse effects are experienced, or C&S report shows resistance or lack of susceptibility of the isolated pathogen to all fluoroquinolones.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

ABSSSI: 14 days. CABP: 10 days.

Prior Authorization Group Description:

BELEODAQ

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

BELSOMRA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

For patients 65 years of age and older: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Rozerem, Silenor 6 mg/day or less, or trazodone. For patients under 65 years of age: Failure of zolpidem or zolpidem CR, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

BENLYSTA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Confirmation that member is positive for autoantibody (e.g., anti-nuclear antibody (ANA), anti-double-stranded DNA (anti-ds-DNA), anti-Smith antigen (anti-Sm), anti-ribonucleoprotein (anti-RNP), anti-Ro/SSA, anti-La/SSB).

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

Currently receiving standard therapy for systemic lupus erythematosus that includes one or more of the following agents, unless all agents are contraindicated: glucocorticoids (e.g., prednisone), antimalarial (e.g., hydroxychloroquine or chloroquine), non-biologic immunosuppressants (e.g., azathioprine, methotrexate, mycophenolate).

Prior Authorization Group Description:

BENZTROPINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Parkinsons disease/Parkinsonism: Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: amantadine, levodopa/carbidopa, entacapone, pramipexole, ropinirole, selegiline.

Prior Authorization Group Description:

BEOVU

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

CONTINUATION OF THERAPY, NEOVASCULAR (WET) AGE-RELATED MACULAR DEGENERATION (AMD): Member is responding positively to therapy as evidenced by one of the following (a, b, c, or d): a) detained neovascularization, b) improvement/stabilization in visual acuity, c) maintenance of corrected visual acuity from prior treatment, or d) supportive findings from optical coherence tomography or fluorescein angiography.

Age Restrictions:

Prescriber Restrictions:

NEOVASCULAR (WET) AMD: Prescribed by or in consultation with an ophthalmologist.

Coverage Duration:

NEOVASCULAR (WET) AMD: Initial: 4 months. Continuation: 6 months.

Other Criteria:

NEOVASCULAR (WET) AMD: Failure of intravitreal bevacizumab, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

BLEOMYCIN

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

BOSULIF

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Confirmation that the member has Philadelphia chromosome positive disease.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

BOTOX

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

CHRONIC MIGRAINE HEADACHE: Persistent history of chronic, debilitating migraine headaches with frequent attacks on more than 15 days per month.

Age Restrictions:

Strabismus or blepharospasm associated with dystonia: 12 years of age or older.

Prescriber Restrictions:

Chronic migraine headache: Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Other Criteria:

Chronic migraine headache: Failure of prophylactic treatment with ONE of the following, unless contraindicated or clinically significant adverse effects are experienced: divalproex, topiramate, timolol or propranolol AND Failure of abortive therapy with ONE of the following, unless contraindicated or clinically significant adverse effects are experienced: sumatriptan, rizatriptan, zolmitriptan, naratriptan, almotriptan, frovatriptan, Relpax, ergotamine/caffeine or dihydroergotamine.

Prior Authorization Group Description:

BRAFTOVI

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Positive for BRAF V600E or V600K mutation.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

MELANOMA: Prescribed in combination with Mektovi. COLON CANCER, RECTAL CANCER: Prescribed in combination with either Erbitux or Vectibix.

Prior Authorization Group Description:

BRIVIACT

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of two of the following generic antiepileptic drugs, unless contraindicated or clinically significant adverse effects are experienced: lamotrigine, topiramate, oxcarbazepine, carbamazepine, phenytoin, valproic acid or divalproex sodium.

Prior Authorization Group Description:

BRUKINSA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

MANTLE CELL LYMPHOMA: Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Other Criteria:

MANTLE CELL LYMPHOMA: Previously received at least one prior therapy (e.g., rituximab-containing regimen).

Prior Authorization Group Description:

C1 ESTERASE INHIBITOR

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Member is not using the requested product in combination with another FDA-approved product for the same indication (e.g., using both Cinryze and Haegarda for long-term prophylaxis of HAE attacks).

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an immunologist, allergist, or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

CABLIVI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Prescribed in combination with plasma exchange therapy. Prescribed in combination with immunosuppressive therapy (i.e., glucocorticoids, rituximab). CONTINUATION OF THERAPY: Member has received no more than 58 days of Cablivi therapy after completion of plasma exchange therapy AND member meets one of the following (a or b): a) If request is for a new treatment cycle, member has experienced no more than two recurrences while taking Cablivi and Cablivi is prescribed in combination with plasma exchange and immunosuppressive therapy (i.e., glucocorticoids, rituximab) OR b) If request is for treatment extension, member is responding positively to therapy as evidenced by, including but not limited to, improvement in any of the following parameters: increase in platelet counts, reduction in neurological symptoms, or improvements in organ-damage markers (lactate dehydrogenase, cardiac troponin I, and serum creatinine).

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a hematologist.

Coverage Duration:

Initial: 60 days. Reauthorization: 58 days post plasma-exchange.

Prior Authorization Group Description:

CABOMETYX

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

NON-SMALL CELL LUNG CANCER: Confirmation of an RET gene rearrangement.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with oncologist.

Coverage Duration:

12 months.

Other Criteria:

HEPATOCELLULAR CARCINOMA: Failure of Nexavar, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

CALQUENCE

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

CLL/SLL: If disease is refractory to Imbruvica, member does not have BTK C481S mutations.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Other Criteria:

MCL: Previously received at least one prior therapy (e.g., rituximab-containing regimen).

Prior Authorization Group Description:

CAPLYTA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

SCHIZOPHRENIA: Failure of two of the following atypical antipsychotics, unless contraindicated or clinically significant adverse effects are experienced: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole.

Prior Authorization Group Description:

CAPRELSA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

NON-SMALL CELL LUNG CANCER: Confirmation of an RET gene rearrangement.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

DIFFERENTIATED THYROID CARCINOMA: Failure of Lenvima or Nexavar, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

CARISOPRODOL/ASPIRIN/CODEINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

CAYSTON

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

CERDELGA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Extensive metabolizer (EM) or intermediate metabolizer (IM) taking a strong or moderate CYP2D6 inhibitor concomitantly with a strong or moderate CYP3A inhibitor AND IMs or poor metabolizer (PM) taking a strong CYP3A inhibitor.

Required Medical Information:

An FDA-cleared genotyping test has determined that this patient is a CYP2D6 extensive metabolizer (EM), intermediate metabolizer (IM), or poor metabolizer (PM).

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

CEREZYME

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Type 3 Gaucher disease.

Exclusion Criteria:

Required Medical Information:

Confirmation of at least one of the following conditions resulting from Gaucher disease: anemia, thrombocytopenia, bone disease, hepatomegaly or splenomegaly.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

CHLORZOXAZONE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

CHORIONIC GONADOTROPIN

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Induction of ovulation and pregnancy in the anovulatory, infertile woman in whom the cause of anovulation is secondary and not due to primary ovarian failure. Treatment of obesity.

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

CIALIS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Member is on concomitant nitrates (e.g., Nitrodur, Nitrobid, Nitrostat, Isordil, Ismo) or guanylate cyclase stimulators (e.g., Adempas (riociguat)).

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

BENIGN PROSTATIC HYPERPLASIA: Failure of ONE alpha blocker (e.g., terazosin, doxazosin, tamsulosin, alfuzosin, Rapaflo) and ONE 5-alpha reductase inhibitor (finasteride, dutasteride/tamsulosin, or dutasteride), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

CIMZIA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

CROHN'S DISEASE: Prescribed by or in consultation with a gastroenterologist. PSORIATIC ARTHRITIS, PLAQUE PSORIASIS: Prescribed by or in consultation with a rheumatologist or dermatologist. RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS, NON-RADIOGRAPHIC AXIAL SPONDYLOARTHRITIS: Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. PLAQUE PSORIASIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine, or acitretin.

Prior Authorization Group Description:

CINQAIR

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Blood eosinophil count of greater than or equal to 400 cells/mcL within the past 3 months.

Age Restrictions:

18 years of age or older.

Prescriber Restrictions:

Prescribed by or in consultation with an allergist, pulmonologist, or immunologist.

Coverage Duration:

12 months.

Other Criteria:

Prescribed in combination with ONE inhaled corticosteroid (e.g., beclomethasone, budesonide, flunisolide, fluticasone, mometasone, ciclesonide), unless contraindicated or clinically significant adverse effects are experienced. AND Prescribed in combination with ONE long-acting beta-agonist (e.g., salmeterol, formoterol, vilanterol), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

CLADRIBINE

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

CLOMIPRAMINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Autistic disorder.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of one selective serotonin reuptake inhibitor (e.g., fluoxetine, fluvoxamine, sertraline), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

COMETRIQ

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

NON-SMALL CELL LUNG CANCER: Confirmation of an RET gene rearrangement.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

DIFFERENTIATED THYROID CARCINOMA: Failure of Lenvima or Nexavar unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

COPIKTRA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

COSENTYX

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

PSORIATIC ARTHRITIS, PLAQUE PSORIASIS: Prescribed by or in consultation with a rheumatologist or dermatologist. ANKYLOSING SPONDYLITIS: Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

PLAQUE PSORIASIS: Failure of ONE of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine, or acitretin. PSORIATIC ARTHRITIS: Failure of one TNF inhibitor (e.g., Humira, Enbrel, Cimzia, Simponi, Simponi Aria, Remicade, Inflectra), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

COTELLIC

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Patients with wild-type BRAF melanoma.

Required Medical Information:

Disease is positive for the BRAF V600E or V600K mutation.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

Prescribed in combination with Zelboraf.

Prior Authorization Group Description:

CRINONE

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Progesterone supplementation or replacement as part of an Assisted Reproductive Technology (ART) treatment for infertile women with progesterone deficiency.

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Crinone 8%: Member has failed to respond or has had an inadequate response to Crinone 4%.

Prior Authorization Group Description:

CROFELEMER

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Patient is on anti-retroviral therapy and failure of loperamide or diphenoxylate/atropine, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

CRYSVITA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

X-LINKED HYPOPHOSPHATEMIA: DNA testing results confirm the presence of mutations in the PHEX gene or documentation of elevated serum fibroblast growth factor 23 (FGF23) levels. Current (within the last 30 days) serum phosphorus level is below the reference range for age and gender. CONTINUATION OF THERAPY: Member meets all approval criteria and has had an increase in serum phosphorus level from baseline and/or maintenance within the normal range for age and gender, while on Crysvita therapy.

Age Restrictions:

At least 6 months of age.

Prescriber Restrictions:

X-LINKED HYPOPHOSPHATEMIA: Prescribed by or in consultation with an endocrinologist or metabolic disease specialist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

CYCLOBENZAPRINE HCL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

CYTARABINE

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Other Criteria:

For acute non-lymphocytic leukemia: use in combination with other approved anti-cancer drugs.

Prior Authorization Group Description:

DAURISMO

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age 75 years or greater, OR medical justification supports inability to use intensive induction chemotherapy OR member responded to then relapsed after Daurismo induction therapy 12 or more months ago. Prescribed in combination with low-dose cytarabine.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

DICLOFENAC GEL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

6 months.

Prior Authorization Group Description:

DIPYRIDAMOLE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

DISOPYRAMIDE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

DOPTELET

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

THROMBOCYTOPENIA WITH CHRONIC LIVER DISEASE: Recent (within the past 14 days) platelet count is less than 50 x 10^9/L. Member is scheduled to undergo a medical or dental procedure within the next 30 days.

Age Restrictions:

Prescriber Restrictions:

THROMBOCYTOPENIA WITH CHRONIC LIVER DISEASE: Prescribed by or in consultation with a hematologist, hepatologist, or gastroenterologist. CHRONIC IMMUNE THROMBOCYTOPENIA (ITP): Prescribed by or in consultation with a hematologist.

Coverage Duration:

THROMBOCYTOPENIA WITH CHRONIC LIVER DISEASE: 4 weeks. CHRONIC ITP: 12 months.

Other Criteria:

THROMBOCYTOPENIA WITH CHRONIC LIVER DISEASE: For members with platelet count less than 40 x 10^9/L, failure of Mulpleta unless contraindicated or clinically significant adverse effects are experienced. CHRONIC ITP: Failure of a corticosteroid (e.g., prednisone, methylprednisolone, or dexamethasone), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

DOXEPIN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

DOXEPIN CREAM

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

4 weeks.

Other Criteria:

Failure of two topical corticosteroids (e.g., amcinonide, fluticasone propionate, triamcinolone acetonide, betamethasone valerate, fluocinolone acetonide, hydrocortisone butyrate, mometasone furoate, desoximetasone, fluocinonide or betamethasone dipropionate), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

DUEXIS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: pantoprazole, lansoprazole or omeprazole AND Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: ibuprofen, diclofenac sodium or potassium, etodolac, fenoprofen, ketoprofen, meloxicam, naproxen, oxaprozin, piroxicam, salsalate, sulindac, tolmetin.

Prior Authorization Group Description:

ELIDEL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of two medium to high potency topical corticosteroids (e.g., amcinonide, fluticasone propionate, triamcinolone acetonide, betamethasone valerate, fluocinolone acetonide, hydrocortisone butyrate, mometasone furoate, desoximetasone, fluocinonide or betamethasone dipropionate), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

EMEND 40 MG

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

4 weeks.

Prior Authorization Group Description:

EMFLAZA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Diagnosis of Duchenne muscular dystrophy (DMD) confirmed by one of the following: Genetic testing (e.g., dystrophin deletion or duplication mutation found) OR if genetic studies are negative (i.e., no mutation identified), positive muscle biopsy (e.g., absence of dystrophin protein).

Age Restrictions:

2 years of age or older.

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of prednisone, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

EMGALITY

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

MIGRAINE PROPHYLAXIS: One of the following (a or b): a) Member experiences 4 or more migraine days per month for at least 3 months, or b) Diagnosis of chronic migraine and Emgality is prescribed for prophylaxis. EPISODIC CLUSTER HEADACHE: Member has had at least 2 cluster headache attack periods which lasted for 1 year or less each and were separated by at least 3 months. CONTINUATION OF THERAPY, MIGRAINE PROPHYLAXIS: Member has experienced and maintained positive response to therapy as evidenced by a reduction in migraine days per month from baseline. CONTINUATION OF THERAPY, EPISODIC CLUSTER HEADACHE: Member has experienced and maintained positive response to therapy as evidenced by a reduction in cluster headache attack frequency. Member has not received more than 12 months of consecutive treatment OR it has been at least 3 months since the member last received Emgality.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist, headache, or pain specialist.

Coverage Duration:

Initial: 3 months. Reauthorizations: 6 months.

Other Criteria:

MIGRAINE PROPHYLAXIS: Failure of two of the following oral migraine preventative therapies, each from different therapeutic classes, unless contraindicated or clinically significant adverse effects are experienced: antiepileptic drugs (e.g., divalproex sodium, sodium valproate, topiramate), beta-blockers (e.g., metoprolol, propranolol, timolol), antidepressants (e.g., amitriptyline, venlafaxine).

Prior Authorization Group Description:

ENBREL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Hidradenitis suppurativa.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

PSORIATIC ARTHRITIS, PLAQUE PSORIASIS: Prescribed by or in consultation with a rheumatologist or dermatologist. RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist. HIDRADENITIS SUPPURATIVA: Prescribed by or in consultation with a rheumatologist, dermatologist or gastroenterologist.

Coverage Duration:

12 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. PLAQUE PSORIASIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin.

Prior Authorization Group Description:

ENDARI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Age 5 or older.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

ENTRESTO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Left ventricular ejection fraction less than or equal to 35%.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a cardiologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

ENTYVIO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a gastroenterologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of Humira or Remicade, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

EPCLUSA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Criteria will be applied consistent with current AASLD-IDSA guidance.

Exclusion Criteria:

Required Medical Information:

Refer to the document, "Recommendations for Testing, Managing, and Treating Hepatitis C," by AASLD-IDSA available at http://www.hcvguidelines.org for drug regimen and duration of treatment based on genotype, treatment status, previous drug regimens used, past medical history and comorbidities.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a gastroenterologist, hepatologist or infectious disease physician.

Coverage Duration:

12 to 24 weeks based on cirrhosis status, genotype, prior treatment, or ribavirin eligibility.

Prior Authorization Group Description:

EPIDIOLEX

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

LENNOX-GASTAUT SYNDROME: will be used as adjunctive therapy with other antiepileptic drugs (e.g., Banzel, clobazam, clonazepam, felbamate, lamotrigine, or topiramate).

Age Restrictions:

Age greater than or equal to 2 years.

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Other Criteria:

LENNOX-GASTAUT SYNDROME: Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: Banzel, clobazam, clonazepam, felbamate, lamotrigine, topiramate.

Prior Authorization Group Description:

EPOETIN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Anemia due to myelodyspastic syndrome. Anemia associated with myelofibrosis. Anemia secondary to combination ribavirin and interferon-alfa therapy in patients infected with hepatitis C virus.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

ERGOLOID MESYLATES

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Alzheimer's dementia: Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: donepezil, memantine, rivastigmine or galantamine.

Prior Authorization Group Description:

ERLEADA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

PROSTATE CANCER: Concurrent use of a gonadotropin-releasing hormone (GnRH) analog or past bilateral orchiectomy. Disease is characterized as one of the following (a or b): a) not metastatic and castration-resistant, OR b) metastatic and castration-sensitive.

Age Restrictions:

Prescriber Restrictions:

PROSTATE CANCER: Prescribed by or in consultation with an oncologist or urologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

ESBRIET

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

ESTROGENS(Duavee, Divigel, Climara, Elestrin, Mimvey, Prempro, Amabelz, Combipatch, Climara Pro, Evamist, Estrace, Activella, Menostar, Vivelle-Dot, Femhrt, Mimvey Lo, Alora, Lopreeza, Premarin, Premphase)

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Atrophic Vaginitis and Kraurosis Vulvae: Failure to one of the following, unless contraindicated or clinically significant adverse effects are experienced: Estradiol vaginal tablet, Femring or Premarin vaginal cream.

Prior Authorization Group Description:

EVZIO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of Narcan (naloxone nasal spray), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

EXONDYS 51

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Duchenne muscular dystrophy with mutation amenable to exon 51 skipping confirmed by genetic testing.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

6 months.

Other Criteria:

Currently stable on an oral corticosteroid regimen (e.g., prednisone), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

EYLEA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an ophthalmologist.

Coverage Duration:

12 months.

Other Criteria:

For all indications except for DME in members with baseline visual acuity worse than 20/50: Failure of bevacizumab unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

FARYDAK

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a hematologist or oncologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of two prior regimens, including bortezomib and an immunomodulatory agent (e.g., dexamethasone), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

FASENRA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Blood eosinophil count of greater than or equal to 150 cells/mcL within the past 3 months.

Age Restrictions:

12 years of age or older.

Prescriber Restrictions:

Prescribed by or in consultation with an allergist, pulmonologist, or immunologist.

Coverage Duration:

12 months.

Other Criteria:

Prescribed in combination with ONE inhaled corticosteroid (e.g., beclomethasone, budesonide, flunisolide, fluticasone, mometasone, ciclesonide), unless contraindicated or clinically significant adverse effects are experienced AND Prescribed in combination with ONE long-acting beta-agonist (e.g., salmeterol, formoterol, vilanterol), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

FENTORA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Age 18 or greater.

Prescriber Restrictions:

Coverage Duration:

Through the end of the Plan contract year.

Other Criteria:

Patient is already taking and is tolerant to around-the-clock opioid therapy. Patients are considered opioid tolerant when taking another opioid daily for a week or longer (for example, at least 60 mg of oral morphine per day or an equianalgesic dose of another opioid).

Prior Authorization Group Description:

FERRIPROX

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of deferoxamine, Exjade or Jadenu, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

FINTEPLA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

DRAVET SYNDROME: Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

FIORICET WITH CODEINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of naproxen and ibuprofen, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

FIORINAL WITH CODEINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of naproxen and ibuprofen, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

FIRAZYR

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Member is not using Firazyr in combination with another FDA-approved product for treatment of acute HAE attacks (e.g., Berinert, Ruconest, Kalbitor).

Age Restrictions:

Age 18 or greater.

Prescriber Restrictions:

Prescribed by or in consultation with an immunologist, allergist, or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

FIRDAPSE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Confirmation of a baseline clinical muscle strength assessment (examples may include but are not limited to the Quantitative Myasthenia Gravis (QMG) score, triple-timed up-and-go test (3TUG), Timed 25-foot Walk test (T25FW)). CONTINUATION OF THERAPY: Member is responding positively to therapy as evidenced by clinical muscle strength assessments.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

FLECTOR

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Cancer pain.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Acute Pain: 4 weeks. Cancer pain: Through the end of the Plan contract year.

Prior Authorization Group Description:

FLUOROURACIL

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

FORTEO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Total duration of therapy on parathyroid hormone (PTH) analogs (e.g., Tymlos, Forteo) has not exceeded 2 years.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Member meets one of the following (a, b, or c): a) Failure of a bisphosphonate (e.g., alendronate) unless contraindicated or clinically significant adverse effects are experienced. OR b) Bone mineral density T-score at hip or spine is -3.5 or less. OR c) Bone mineral density T-score at hip or spine is -2.5 or less with a history of major osteoporotic fracture of the hip, spine, forearm, wrist, or humerus.

Prior Authorization Group Description:

GALAFOLD

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Amenable GLA variants (mutations) associated with benign phenotypes (i.e., phenotypes known not to cause Fabry disease), including the following GLA mutation: c.937G to T, (p.(D313Y)).

Required Medical Information:

Presence of at least one amenable GLA variant (mutation) as confirmed by one of the following resources: Galafold Prescribing Information brochure (package insert - Section 12, Table 2), Amicus Fabry GLA Gene Variant Search Tool: http://www.fabrygenevariantsearch.com/hcp, or Amicus Medical Information at 1-877-4AMICUS or medinfousa@amicusrx.com.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a clinical geneticist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

GANCICLOVIR

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

GATTEX

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Member has been dependent on parenteral nutrition or other intravenous support for at least 12 months. CONTINUATION OF THERAPY: Requirement for parenteral nutrition or other intravenous support has decreased since initiation of Gattex therapy.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a gastroenterologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

GAVRETO

Pending CMS Review

Prior Authorization Group Description:

GILENYA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Baseline QTc interval greater than or equal to 500 msec.

Required Medical Information:

Age Restrictions:

10 years of age or older.

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

GILOTRIF

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

NON-SMALL CELL LUNG CANCER: Member meets one of the following (a or b): a) disease is positive for a sensitizing EGFR mutation (e.g., exon 19 deletion or insertion, exon 21 point mutation (L858R, L861Q), exon 18 point mutation (G719X), exon 20 point mutation (S768I)), OR b) disease is squamous cell carcinoma histology with progression after platinum-based chemotherapy (e.g., cisplatin, carboplatin).

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

GLATIRAMER

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

GLIMEPIRIDE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: glipizide or glipizide/metformin combination product.

Prior Authorization Group Description:

GLYBURIDE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: glipizide or glipizide/metformin combination product.

Prior Authorization Group Description:

GLYBURIDE/METFORMIN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: glipizide or glipizide/metformin combination product.

Prior Authorization Group Description:

GRANIX

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Myelodysplastic syndrome.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

HARVONI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Criteria will be applied consistent with current AASLD-IDSA guidance.

Exclusion Criteria:

Required Medical Information:

Refer to the document, "Recommendations for Testing, Managing, and Treating Hepatitis C," by AASLD-IDSA available at http://www.hcvguidelines.org for drug regimen and duration of treatment based on genotype, treatment status, previous drug regimens used, past medical history and comorbidities.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a gastroenterologist, hepatologist or infectious disease physician.

Coverage Duration:

8 to 24 weeks based on cirrhosis status, genotype, prior treatment, or ribavirin eligibility.

Prior Authorization Group Description:

HERCEPTIN

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Disease is human epidermal growth factor receptor (HER2) positive.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

GASTRIC AND ESOPHAGEAL CANCER: prescribed in combination with systemic chemotherapy (e.g., cisplatin and either capecitabine or 5-fluorouracil). ENDOMETRIAL CARCINOMA: prescribed in combination with carboplatin and paclitaxel.

Prior Authorization Group Description:

HETLIOZ

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

HUMAN GROWTH HORMONE

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

CHILDREN AND ADOLESCENTS WITH GROWTH HORMONE DEFICIENCY, SHOX DEFICIENCY IN CHILDREN: Baseline height must be greater than 2 standard deviations below the mean for gender and age. Growth rate is such that the member is unlikely to attain an adult height in the normal range - 59 inches for girls and 63 inches for boys. TURNER SYNDROME: Confirmed by karyotype. PRADER-WILLI or NOONAN SYNDROME: Baseline height must be less than the 5th percentile for gender and age OR 2 or more standard deviations below the mean measured paternal height. Growth rate is such that the member is unlikely to attain an adult height in the normal range - 59 inches for girls and 63 inches for boys.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Adult Growth Hormone Deficiency: 12 months. HIV Wasting or Cachexia, Children: 6 months.

Other Criteria:

HIV Wasting or Cachexia: Member is being treated with concomitant antiretroviral therapy.

Prior Authorization Group Description:

HUMIRA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

For the following indications, prescribed by or in consultation with: PSORIATIC ARTHRITIS, PLAQUE PSORIASIS - rheumatologist or dermatologist. CROHN'S DISEASE, ULCERATIVE COLITIS gastroenterologist. RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS - rheumatologist. HIDRADENITIS SUPPURATIVA - rheumatologist, dermatologist or gastroenterologist. UVEITIS - ophthalmologist or rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. PLAQUE PSORIASIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin.

Prior Authorization Group Description:

HYDROCODONE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

3 months initial for non-malignant pain then 12 months. 12 months for cancer pain.

Other Criteria:

Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: MS Contin, Kadian, Duragesic, Opana ER, Avinza or Oxycontin.

Prior Authorization Group Description:

HYDROXYZINE HCL INJECTION

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

HYDROXYZINE HCL ORAL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Pruritus: Failure of one of the following topical agents, unless contraindicated or clinically significant adverse effects are experienced: betamethasone, hydrocortisone, triamcinolone, fluticasone, clobetasol, fluocinonide or fluocinolone. Anxiety: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: venlafaxine, buspirone, duloxetine or escitalopram. All other FDA approved indications: Patient is continuing on this medication without adverse effects.

Prior Authorization Group Description:

HYDROXYZINE PAMOATE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Pruritus: Failure of one of the following topical agents, unless contraindicated or clinically significant adverse effects are experienced: betamethasone, hydrocortisone, triamcinolone, fluticasone, clobetasol, fluocinonide or fluocinolone. Anxiety: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: venlafaxine, buspirone, duloxetine or escitalopram. All other FDA approved indications: Patient is continuing on this medication without adverse effects.

Prior Authorization Group Description:

ICLUSIG

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

ACUTE LYMPHOBLASTIC LEUKEMIA (ALL): disease is Philadelphia chromosome positive (Ph+).

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

IDHIFA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Presence of an isocitrate dehydrogenase-2 (IDH2) mutation.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Other Criteria:

For age less than 60 years, disease has relapsed or is refractory following treatment with a first line chemotherapy regimen (e.g., cytarabine, idarubicin, daunorubicin, Vyxeos, cladribine, Rydapt, Mylotarg).

Prior Authorization Group Description:

ILARIS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Acute gouty arthritis.

Exclusion Criteria:

Required Medical Information:

Confirmation of current weight.

Age Restrictions:

Cryopyrin-Associated Periodic Syndromes: 4 years and older. All other covered indications: 2 years and older.

Prescriber Restrictions:

SYSTEMIC JUVENILE IDIOPATHIC ARTHRITIS: Prescribed by or in consultation with a dermatologist, rheumatologist, or gastroenterologist. ALL OTHER COVERED INDICATIONS: Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

ILUMYA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a rheumatologist or dermatologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin. Failure of one of the following unless contraindicated or clinically significant adverse effects are experienced: Cosentyx, Humira, Inflectra, Remicade, Stelara, Taltz, and Tremfya.

Prior Authorization Group Description:

IMATINIB

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

CHRONIC MYELOID LEUKEMIA (CML), ACUTE LYMPHOBLASTIC LEUKEMIA (ALL): disease is Philadelphia chromosome positive. CHRONIC MYELOMONOCYTIC LEUKEMIA: disease is positive for a platelet-derived growth factor receptor (PDGFR) mutation or a 5q3133 mutation. MYELODYSPLASTIC/MYELOPROLIFERATIVE DISEASES: disease is positive for a PDGFR mutation. AGGRESSIVE SYSTEMIC MASTOCYTOSIS: disease is D816V c-Kit mutation negative or c-Kit mutational status is unknown. MELANOMA: disease is KIT-positive. PIGMENTED VILLONODULAR SYNOVITIS/TENOSYNOVIAL GIANT CELL TUMOR: Disease is associated with severe morbidity or functional limitations and is not amenable to improvement with surgery.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

IMBRUVICA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

CHRONIC GRAFT-VERSUS-HOST DISEASE: Prescribed by or in consultation with an oncologist, hematologist, or bone marrow transplant specialist. ALL OTHER INDICATIONS: Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

IMIPRAMINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Depression: Failure of one of the following generic antidepressants, unless contraindicated or clinically significant adverse effects are experienced: bupropion, bupropion SR, bupropion XL, citalopram, desvenlafaxine succinate, duloxetine, escitalopram, fluoxetine, mirtazapine, sertraline, venlafaxine or venlafaxine XR.

Prior Authorization Group Description:

INDOMETHACIN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of naproxen and sulindac, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

INFLECTRA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Psoriatic Arthritis/Plaque Psoriasis: Prescribed by or in consultation with a rheumatologist or dermatologist. Crohn's Disease/Ulcerative Colitis: Prescribed by or in consultation with a gastroenterologist. Rheumatoid Arthritis/Ankylosing Spondylitis: Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

Rheumatoid Arthritis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. Plaque Psoriasis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin.

Prior Authorization Group Description:

INGREZZA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Development of tardive dyskinesia is secondary to a centrally acting dopamine receptor blocking agent (neuroleptic) (e.g., first- or second-generation antipsychotics such as chlorpromazine or aripiprazole, antiemetics such as promethazine or metoclopramide, the tri-cyclic antidepressant amoxapine).

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a psychiatrist or neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

INLYTA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

RENAL CELL CARCINOMA WITH CLEAR CELL HISTOLOGY: One of the following (a or b): a) prescribed concurrently with Keytruda or Bavencio, OR b) prescribed as monotherapy after failure of one prior therapy (e.g., Votrient, Sutent), unless contraindicated or clinically significant adverse effects are experienced. DIFFERENTIATED THYROID CARCINOMA: Failure of Lenvima or Nexavar unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

INQOVI

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

INREBIC

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

MYELOFIBROSIS: Confirmation of a recent (within the last 30 days) thiamine level of 70 nmol/L (3 mcg/dL) or greater. Confirmation of a recent (within the last 30 days) platelet count of 50,000/mcL or greater.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

INTERFERON BETA-1A

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

INTERFERON BETA-1B

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

INTUNIV

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Attention Deficit Hyperactivity Disorder: Failure of both of the following, unless contraindicated or clinically significant adverse effects are experienced: amphetamine-based stimulant and methylphenidate based-stimulant.

Prior Authorization Group Description:

JAKAFI

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

STEROID REFRACTORY GRAFT-VERSUS-HOST DISEASE: Member has history of bone marrow or stem cell transplant.

Age Restrictions:

STEROID REFRACTORY GRAFT-VERSUS-HOST DISEASE: Age greater than or equal to 12 years.

Prescriber Restrictions:

STEROID REFRACTORY GRAFT-VERSUS-HOST DISEASE: Prescribed by or in consultation with an oncologist, hematologist, or bone marrow transplant specialist. ALL OTHER INDICATIONS: Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Other Criteria:

POLYCYTHEMIA VERA: Failure of hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced. STEROID REFRACTORY GRAFT-VERSUS-HOST DISEASE: Failure of systemic corticosteroids, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

JUBLIA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of terbinafine tablets, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

JUXTAPID

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of Repatha 420 mg, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

JYNARQUE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a nephrologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

KADCYLA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Kadcyla will be used as a single-agent therapy.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

KADIAN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Medical justification as to why patient cannot take an equivalent daily dose of a generically available extended-release morphine sulfate product.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

KALYDECO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Presence of one mutation in the CFTR gene that is responsive to ivacaftor. Confirmation that a homozygous F508del mutation in the CFTR gene is not present.

Age Restrictions:

6 months of age or older.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

KAZANO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of Tradjenta or Jentadueto, unless contraindicated or clinically significant adverse effects are experienced AND Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Januvia, Janumet, Janumet XR.

Prior Authorization Group Description:

KERYDIN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of terbinafine tablets, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

KETOROLAC TROMETHAMINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Patients with active peptic ulcer disease. Advanced renal impairment or at risk for renal failure due to volume depletion. Suspected or confirmed cerebrovascular bleeding, hemorrhagic diathesis, incomplete hemostasis and those at high risk for bleeding. Patient currently receiving aspirin or NSAIDs (non-steroidal anti-inflammatory drugs). Patient currently receiving probenecid or pentoxifylline.

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

5 days.

Prior Authorization Group Description:

KEVEYIS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

KEVZARA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin.

Prior Authorization Group Description:

KEYTRUDA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

CLASSICAL HODGKIN LYMPHOMA, PRIMARY MEDIASTINAL LARGE B-CELL LYMPHOMA: Prescribed by or in consultation with an oncologist or hematologist. UROTHELIAL CARCINOMA: Prescribed by or in consultation with an oncologist or urologist. ALL OTHER INDICATIONS: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

KINERET

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

RHEUMATOID ARTHRITIS: Prescribed by or in consultation with a rheumatologist. NEONATAL-ONSET MULTISYSTEM INFLAMMATORY DISEASE (NOMID): Prescribed by or in consultation with a rheumatologist, neonatologist, or pediatrician.

Coverage Duration:

12 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin AND Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Enbrel, Humira, Remicade, Cimzia, Simponi or Simponi Aria.

Prior Authorization Group Description:

KISQALI(Kisqali Femara Co-Pack, Kisqali)

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Breast cancer is hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, and advanced or metastatic.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

For Kisqali: Prescribed in combination with an aromatase inhibitor (e.g., letrozole, anastrozole, or exemestane), fulvestrant, or tamoxifen. If prescribed in combination with tamoxifen: Medical justification supports need to use tamoxifen over an aromatase inhibitor or fulvestrant. For men receiving an aromatase inhibitor: Prescribed in combination with an agent that suppresses testicular steroidogenesis (e.g., gonadotropin-releasing hormone agonists).

Prior Authorization Group Description:

KOMBIGLYZE XR

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of Tradjenta or Jentadueto, unless contraindicated or clinically significant adverse effects are experienced AND Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Januvia, Janumet, Janumet XR.

Prior Authorization Group Description:

KORLYM

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an endocrinologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

KOSELUGO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

NEUROFIBROMATOSIS TYPE 1 (NF1): Diagnosis is confirmed by positive genetic testing for NF1 or member has at least one diagnostic criteria for NF1 based on the National Institutes of Health Neurofibromatosis 1 Diagnostic Criterion. Complete resection of plexiform neurofibroma (PN) is not considered to be feasible without substantial risk or morbidity (e.g., due to encasement of, or close proximity to, vital structures, invasiveness, or high vascularity of the PN).

Age Restrictions:

Prescriber Restrictions:

NF1: Prescribed by or in consultation with an oncologist or neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

KUVAN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Recent (within 90 days) phenylalanine (Phe) blood level is greater than 360 micromol/L. CONTINUATION OF THERAPY: Confirmation of a reduction in Phe blood levels since initiation of therapy.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a metabolic or genetic disease specialist.

Coverage Duration:

Initial: 3 months. Reauthorization: 12 months.

Prior Authorization Group Description:

LATUDA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of two of the following atypical antipsychotics, unless contraindicated or clinically significant adverse effects are experienced: risperidone, olanzapine, quetiapine, ziprasidone, aripiprazole.

Prior Authorization Group Description:

LAZANDA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Age 18 or greater

Prescriber Restrictions:

Coverage Duration:

Through the end of the Plan contract year.

Other Criteria:

Patient is already taking and is tolerant to around-the-clock opioid therapy. Patients are considered opioid tolerant when taking another opioid daily for a week or longer (for example, at least 60 mg of oral morphine per day or an equianalgesic dose of another opioid).

Prior Authorization Group Description:

LEMTRADA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of TWO of the following, unless contraindicated or clinically significant adverse effects are experienced: Aubagio, Tecfidera, Gilenya, Avonex, Betaseron, Plegridy, glatiramer, Copaxone, Glatopa, Extavia or Rebif.

Prior Authorization Group Description:

LENVIMA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

RENAL CELL CARCINOMA: Prescribed in combination with Afinitor AND if histology is clear cell or unknown, failure of a regimen consisting of or including one of the following drugs unless contraindicated or clinically significant adverse effects are experienced: Avastin, Cabometyx, Inlyta, Nexavar, Opdivo, Proleukin, Sutent, Tarceva, Torisel, Votrient, or Yervoy. MEDULLARY THYROID CARCINOMA: Failure of Cometriq or Caprelsa unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

LEUKINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

AML following induction therapy, Mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation, Following autologous peripheral blood progenitor cell or bone marrow transplantation (BMT) in members with NHL, ALL, HL for acceleration of myeloid reconstitution, Following allogeneic BMT for acceleration of myeloid reconstitution, Acute Radiation Syndrome: Failure of Neupogen, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

LIDODERM

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Diabetic neuropathy. Cancer-related neuropathic pain.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

LONSURF

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

COLORECTAL CANCER: Confirmation that the patient does or does not have the RAS (KRAS or NRAS) wild type gene. GASTRIC CANCER, GASTROESOPHAGEAL ADENOCARCINOMA: Confirmation that the patient does or does not have a HER2/neu-positive tumor.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

COLORECTAL CANCER: If tumor expresses the RAS wild type gene, failure of Erbitux or Vectibix, unless contraindicated or clinically significant adverse effects are experienced. GASTRIC CANCER, GASTROESOPHAGEAL ADENOCARCINOMA: If tumor is HER2/neu-positive (i.e., HER2-overexpressing), failure of Herceptin, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

LORBRENA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

NON-SMALL CELL LUNG CANCER: Disease is ALK or ROS1 positive.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

NON-SMALL CELL LUNG CANCER: For ALK-positive disease, failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Alecensa, Alunbrig, Xalkori, Zykadia. For ROS1-positive disease, failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Rozlytrek, Xalkori, Zykadia.

Prior Authorization Group Description:

LOTRONEX

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Male patients.

Required Medical Information:

Female patient with irritable bowel symptoms persisting for at least 6 months.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

LUCEMYRA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Diagnosis of opioid dependence (may be limited to physiologic dependence/tolerance) or opioid use disorder. Member is currently or will be undergoing abrupt opioid discontinuation within the next seven days and one of the following: member has taken one or more opioids for at least the last three weeks OR an opioid antagonist (e.g., naltrexone) has been or will be administered after a period of opioid use. Medical justification supports why an opioid taper (e.g., with buprenorphine, methadone or other opioid) cannot be used. Lucemyra has not been prescribed for a prior opioid withdrawal event within the last 30 days or medical justification supports retreatment.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a physician specializing in one of the following areas: emergency medicine/inpatient care, pain management, addiction psychiatry.

Coverage Duration:

14 days per course of treatment.

Prior Authorization Group Description:

LYNPARZA TABLET

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

BREAST CANCER: Mutations in the BRCA genes and confirmation of human epidermal growth factor receptor 2 (HER2)-negative disease. METASTATIC CASTRATION-RESISTANT PROSTATE CANCER: Member does not have a PPP2R2A gene mutation.

Age Restrictions:

Prescriber Restrictions:

PROSTATE CANCER: Prescribed by or in consultation with an oncologist or urologist. ALL OTHER ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

MAVENCLAD

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

CONTINUATION OF THERAPY: Member has not yet received 2 courses (4 cycles) lifetime total.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Other Criteria:

RELAPSING-REMITTING MULTIPLE SCLEROSIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Aubagio, Tecfidera, Gilenya, Avonex, Betaseron, Plegridy, glatiramer, Copaxone, Glatopa, Extavia, or Rebif.

Prior Authorization Group Description:

MAVYRET

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Treatment-experienced patients with both NS3/4A protease inhibitor and NS5A inhibitor.

Required Medical Information:

If cirrhosis is present, confirmation of Child-Pugh A status. Refer to the document, "Recommendations for Testing, Managing, and Treating Hepatitis C," by AASLD-IDSA available at http://www.hcvguidelines.org for drug regimen and duration of treatment based on genotype, treatment status, previous drug regimens used, past medical history and comorbidities.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a gastroenterologist, hepatologist or infectious disease physician.

Coverage Duration:

8 to 16 weeks based on genotype, cirrhosis status, prior treatment regimen.

Other Criteria:

If member has been previously treated with an HCV regimen containing NS5A inhibitor or an NS3/4A protease inhibitor, but not both, member has genotype 1.

Prior Authorization Group Description:

MAYZENT

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

CYP2C9*3/*3 genotype.

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Other Criteria:

RELAPSING-REMITTING MULTIPLE SCLEROSIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Aubagio, Tecfidera, Gilenya, Avonex, Betaseron, Plegridy, glatiramer, Copaxone, Glatopa, Extavia, or Rebif.

Prior Authorization Group Description:

MEGACE

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

BREAST CANCER AND ENDOMETRIAL CANCER: Megestrol acetate is being used for palliative treatment.

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

ANOREXIA AND CACHEXIA ASSOCIATED WITH AIDS: Failure of dronabinol and oxandrolone, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

MEGACE ES

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

BREAST CANCER AND ENDOMETRIAL CANCER: Megestrol acetate is being used for palliative treatment.

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

ANOREXIA AND CACHEXIA ASSOCIATED WITH AIDS: Failure of dronabinol and oxandrolone, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

MEKINIST

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

MELANOMA: Positive for BRAF V600E or V600K mutation. NON-SMALL CELL LUNG CANCER, ANAPLASTIC THYROID CANCER: Positive for BRAF V600E mutation.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

NON-SMALL CELL LUNG CANCER, ANAPLASTIC THYROID CANCER: Prescribed in combination with Tafinlar.

Prior Authorization Group Description:

MEKTOVI

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Positive for BRAF V600E or V600K mutation.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

MELANOMA: Prescribed in combination with Braftovi. COLON CANCER, RECTAL CANCER: Prescribed in combination with Braftovi and either Erbitux or Vectibix.

Prior Authorization Group Description:

MEPROBAMATE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: venlafaxine, buspirone, duloxetine or escitalopram.

Prior Authorization Group Description:

METAXALONE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

METHOCARBAMOL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

METHOTREXATE INJ

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS, RHEUMATOID ARTHRITIS: Prescribed by or in consultation with a rheumatologist. PSORIASIS: Prescribed by or in consultation with a rheumatologist or a dermatologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of generic methotrexate injection, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

METHYLDOPA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: amlodipine/benazepril, benazepril/hydrochlorothiazide, captopril, captopril/hydrochlorothiazide, fosinopril, fosinopril/hydrochlorothiazide, lisinopril, lisinopril/hydrochlorothiazide, quinapril, quinapril/hydrochlorothiazide, losartan, losartan/hydrochlorothiazide, valsartan, valsartan/hydrochlorothiazide, irbesartan, irbesartan/hydrochlorothiazide, candesartan, candesartan/hydrochlorothiazide, carvedilol, labetalol, acebutolol, atenolol, bisoprolol, bisoprolol/hydrochlorothiazide, timolol, nadolol, propranolol, metoprolol, metoprolol, nifedipine SR, amlodipine, nicardipine.

Prior Authorization Group Description:

MIRVASO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Erythema of rosacea with papules or pustules: Failure of topical metronidazole, oral doxycycline or Finacea, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

MOZOBIL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Member is scheduled to receive autologous stem cell transplantation.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Other Criteria:

Prescribed in combination with a granulocyte-colony stimulating factor (G-CSF) (i.e., Neupogen, Zarxio, Granix, or Nivestym).

Prior Authorization Group Description:

MULPLETA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Recent (within the past 14 days) platelet count is less than 50 x $10^9/L$. Member is scheduled to undergo a medical or dental procedure within the next 30 days.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a hematologist, hepatologist, or gastroenterologist.

Coverage Duration:

4 weeks.

Prior Authorization Group Description:

NAMENDA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Vascular dementia.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 59 years and younger. Prior authorization is not required for patients 60 years and older.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

NATPARA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Recent (dated within the last 30 days) serum calcium level is greater than 7.5 mg/dL. Recent (dated within the last 30 days) lab result shows sufficient 25-hydroxyvitamin D stores (at least 50 nmol/L or 20 ng/mL). CONTINUATION OF THERAPY: Maintained on therapy with positive response as evidenced by a recent (dated within the last 90 days) serum calcium level within 8-9 mg/dL.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an endocrinologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of an active form of vitamin D (e.g., calcitriol) unless contraindicated or clinically significant adverse events are experienced.

Prior Authorization Group Description:

NAYZILAM

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Member is currently on a stable regimen of antiepileptic drugs.

Age Restrictions:

Age greater than or equal to 12 years.

Prescriber Restrictions:

Prescribed by or in consultation with neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

NERLYNX

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Human epidermal growth factor receptor 2 (HER2)-positive breast cancer.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

NESINA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of Tradjenta or Jentadueto, unless contraindicated or clinically significant adverse effects are experienced AND Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Januvia, Janumet, Janumet XR.

Prior Authorization Group Description:

NEULASTA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Mobilization of peripheral-blood progenitor cells prior to autologous transplantation. Supportive care post autologous hematopoietic cell transplantation.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

NEUPOGEN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Myelodysplastic syndrome. Neutropenia in patients with HIV/AIDS.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

NINLARO

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Other Criteria:

MULTIPLE MYELOMA: Prescribed in combination with dexamethasone.

Prior Authorization Group Description:

NIVESTYM

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Myelodysplastic syndrome. Neutropenia in patients with HIV/AIDS. Hematopoietic syndrome of acute radiation syndrome.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

NORTHERA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

NUBEQA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Concurrent use of a gonadotropin-releasing hormone (GnRH) analog (e.g., Zoladex, Vantas, leuprolide/Lupron Depot, Eligard, Trelstar, Firmagon) or past bilateral orchiectomy. Disease is not metastatic.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or urologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

NUCALA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

ASTHMA OR EOSINOPHILIC GRANULOMATOSIS WITH POLYANGIITIS: Blood eosinophil count of greater than or equal to 150 cells/mcL within the past 3 months.

Age Restrictions:

ASTHMA: 6 years of age or older.

Prescriber Restrictions:

ASTHMA: Prescribed by or in consultation with an allergist, pulmonologist, or immunologist. EOSINOPHILIC GRANULOMATOSIS WITH POLYANGIITIS: Prescribed by or in consultation with a pulmonologist, immunologist, rheumatologist, or nephrologist.

Coverage Duration:

12 months.

Other Criteria:

ASTHMA: Prescribed in combination with ONE inhaled corticosteroid (e.g., beclomethasone, budesonide, flunisolide, fluticasone, mometasone, ciclesonide), unless contraindicated or clinically significant adverse effects are experienced AND Prescribed in combination with ONE long-acting beta-agonist (e.g., salmeterol, formoterol, vilanterol), unless contraindicated or clinically significant adverse effects are experienced. EOSINOPHILIC GRANULOMATOSIS WITH POLYANGIITIS: Failure of ONE glucocorticoid, unless contraindicated or clinically significant adverse experienced.

Prior Authorization Group Description:

NUEDEXTA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

NUPLAZID

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

NUVIGIL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

NUZYRA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Request is for continuation of therapy initiated in an acute care hospital from which member was discharged OR Culture and sensitivity (C&S) report for the current infection shows isolated pathogen is an organism susceptible to omadacycline, unless provider confirms that obtaining a C&S report is not feasible.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

14 days.

Other Criteria:

For members initiating Nuzyra therapy outside of an acute care hospital, one of the following (a, b, or c): a) If a C&S report is available: Failure of 2 antibiotics to which the isolated pathogen is susceptible (if available) per C&S report, unless all are contraindicated or clinically significant adverse effects are experienced. b) C&S report shows resistance or lack of susceptibility of the isolated pathogen to all antibiotics FDA-approved for member's diagnosis. c) If provider confirms that obtaining a C&S report is not feasible: Failure of 2 antibiotics indicated for member's diagnosis (if available), unless all are contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

OCALIVA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a hepatologist or gastroenterologist.

Coverage Duration:

12 months.

Other Criteria:

Prescribed in combination with ursodeoxycholic acid unless contraindicated or clinically significant adverse effects are experienced to ursodeoxycholic acid.

Prior Authorization Group Description:

OCREVUS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Other Criteria:

RELAPSING-REMITTING MULTIPLE SCLEROSIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Aubagio, Tecfidera, Gilenya, Avonex, Betaseron, Plegridy, glatiramer, Copaxone, Glatopa, Extavia or Rebif.

Prior Authorization Group Description:

ODOMZO

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

OFEV

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

SYSTEMIC SCLEROSIS (SSc) ASSOCIATED INTERSTITIAL LUNG DISEASE: Pulmonary fibrosis on high resolution computed tomography (HRCT). Additional signs of SSc are identified (examples may include but are not limited to skin thickening of the fingers, fingertip lesions, telangiectasia, abnormal nailfold capillaries, Raynaud's phenomenon, pulmonary arterial hypertension, SSc-related autoantibodies - anticentromere, anti-topoisomerase I [anti-Scl-70], anti-RNA polymerase III). CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE WITH A PROGRESSIVE PHENOTYPE: For new starts only: confirmation of both of the following within the past 24 months (a and b): a) pulmonary fibrosis affecting more than 10% of lung volume on HRCT and b) confirmation of one of the following (i or ii): i) a relative decline in the forced vital capacity (FVC) of 10% or more of the predicted value, or ii) a relative decline in the FVC of 5% to less than 10% of the predicted value plus either worsening of respiratory symptoms or an increased extent of fibrosis on HRCT.

Age Restrictions:

Prescriber Restrictions:

SYSTEMIC SCLEROSIS ASSOCIATED INTERSTITIAL LUNG DISEASE, CHRONIC FIBROSING INTERSTITIAL LUNG DISEASE WITH A PROGRESSIVE PHENOTYPE: Prescribed by or in consultation with pulmonologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

OLUMIANT

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following agents, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine, or auranofin. Failure of at least one TNF inhibitor unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

ONGLYZA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of Tradjenta or Jentadueto, unless contraindicated or clinically significant adverse effects are experienced AND Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Januvia, Janumet, Janumet XR.

Prior Authorization Group Description:

ONUREG

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

ACUTE MYELOID LEUKEMIA (AML): Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

End of Plan Year.

Other Criteria:

AML: Medical justification supports inability to use subcutaneous or intravenous azacitidine (e.g., contraindication to excipients).

Prior Authorization Group Description:

OPSUMIT

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

ORALAIR

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Confirmation of a positive skin test or in vitro testing for pollen-specific IgE antibodies for any of the following grass species: Sweet Vernal, Orchard, Perennial Rye, Timothy, and Kentucky Blue Grass.

Age Restrictions:

Age greater than or equal to 5 years and less than or equal to 65 years.

Prescriber Restrictions:

Prescribed by or in consultation with an allergist or immunologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of both of the following, unless contraindicated or clinically significant adverse effects are experienced: oral antihistamines and intranasal corticosteroids.

Prior Authorization Group Description:

ORENCIA CLICKJECT

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

RHEUMATOID ARTHRITIS, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist. PSORIATIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist or dermatologist.

Coverage Duration:

12 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. PSORIATIC ARTHRITIS: Failure of one TNF inhibitor (e.g., Humira, Enbrel, Cimzia, Simponi, Simponi Aria, Remicade, Inflectra), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

ORENCIA IV

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

RHEUMATOID ARTHRITIS, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist. PSORIATIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist or dermatologist.

Coverage Duration:

12 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of Remicade and one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin.

Prior Authorization Group Description:

ORENCIA SC

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

RHEUMATOID ARTHRITIS, POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist. PSORIATIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist or dermatologist.

Coverage Duration:

12 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. PSORIATIC ARTHRITIS: Failure of one TNF inhibitor (e.g., Humira, Enbrel, Cimzia, Simponi, Simponi Aria, Remicade, Inflectra), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

ORENITRAM

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

ORILISSA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

For 200 mg twice daily requests, members with osteoporosis.

Required Medical Information:

Continuation of therapy: improvement in dysmenorrhea, dyspareunia, pelvic pain/induration/tenderness, or size of endometrial lesions. Total duration of therapy has not exceeded 6 months for 200 mg twice daily or 24 months for 150 mg once daily dosing.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a gynecologist.

Coverage Duration:

200 mg twice daily: 6 months. 150 mg once daily: 12 months.

Other Criteria:

Failure of ONE non-steroidal anti-inflammatory drug (e.g., ibuprofen, naproxen, fenoprofen, ketoprofen, mefenamic acid, meclofenamate, indomethacin, tolmetin, diclofenac, etodolac, diflunisal, meloxicam, piroxicam) or ONE progestin-containing agent (e.g., norethindrone, ethinyl estradiol with (desogestrel, ethynodiol diacetate, drospirenone, etonogestrel, levonorgestrel, norelgestromin, norethindrone, norgestimate, or norgestrel), estradiol valerate/dienogest, mestranol/norethindrone, depot injectable medroxyprogesterone acetate), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

ORKAMBI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Presence of homozygous F508del mutation in an FDA-cleared cystic fibrosis mutation test.

Age Restrictions:

2 years of age or older.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

ORPHENADRINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

OSENI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of Tradjenta or Jentadueto, unless contraindicated or clinically significant adverse effects are experienced AND Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Januvia, Janumet, Janumet XR.

Prior Authorization Group Description:

OTEZLA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

PSORIATIC ARTHRITIS, PLAQUE PSORIASIS, ORAL ULCERS ASSOCIATED WITH BEHCET'S DISEASE: Prescribed by or in consultation with a rheumatologist or dermatologist.

Coverage Duration:

12 months.

Other Criteria:

PLAQUE PSORIASIS: Failure to ONE of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin.

Prior Authorization Group Description:

OXBRYTA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

SICKLE CELL DISEASE: Disease is associated with one of the following genotypes: Homozygous hemoglobin S, Hemoglobin S beta 0-thalassemia, Hemoglobin S beta+ thalassemia, Hemoglobin SC. Member has a hemoglobin level between 5.5 and 10.5 g/dL. Member meets one of the following (a or b): a) Member experienced at least 1 vaso-occlusive crisis (VOC) within the past 6 months while on hydroxyurea, OR b) Member has intolerance or contraindication to hydroxyurea and has experienced at least 1 VOC within the past 12 months. Oxbryta is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced. Oxbryta is not prescribed concurrently with Adakveo. CONTINUATION OF THERAPY, SICKLE CELL DISEASE: Member is responding positively to therapy as evidenced by an increase in Hb level from baseline of at least 1 g/dL. Oxbryta is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced. Oxbryta is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse of the form baseline of at least 1 g/dL. Oxbryta is prescribed concurrently with hydroxyurea, unless contraindicated or clinically significant adverse.

Age Restrictions:

SICKLE CELL DISEASE: Age greater than or equal to 12 years.

Prescriber Restrictions:

SICKLE CELL DISEASE: Prescribed by or in consultation with a hematologist.

Coverage Duration:

6 months.

Other Criteria:

SICKLE CELL DISEASE: Failure of L-glutamine, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

OXERVATE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an ophthalmologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

PALYNZIQ

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Recent (within 90 days) phenylalanine (Phe) blood level is greater than 600 micromol/L. CONTINUATION OF THERAPY: Positive response as evidenced by one of the following (a, b, or c): a) Blood Phe level has decreased by at least 20% from pre-treatment baseline, b) Blood Phe level is less than or equal to 600 micromol/L, c) Member has been using 20 mg per day for at least 6 months, but a dose titration to 40 mg per day is being requested after failure to meet therapeutic targets (a or b above).

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an endocrinologist, metabolic disease specialist, or genetic disease specialist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

PEMAZYRE

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

PENNSAID

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of ONE oral non-steroidal anti-inflammatory drug (e.g., ibuprofen, naproxen, fenoprofen, ketoprofen, mefenamic acid, meclofenamate, indomethacin, tolmetin, diclofenac, etodolac, diflunisal, meloxicam, piroxicam), unless all are contraindicated or clinically significant adverse effects are experienced. Failure of either diclofenac 1.5% topical solution or diclofenac 1% topical gel, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

PERSERIS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Member meets one of the following (a or b): a) therapy was initiated in an inpatient setting during a recent (within 60 days) hospital admission, OR b) failure of TWO of the following atypical antipsychotics, unless contraindicated or clinically significant adverse effects are experienced: aripiprazole, ziprasidone, quetiapine, olanzapine, risperidone, asenapine, iloperidone, paliperidone.

Prior Authorization Group Description:

PHENOBARBITAL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Partial seizures: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: carbamazepine, phenytoin, topiramate, tiagabine, levetiracetam, gabapentin, lamotrigine, oxcarbazepine, primidone or divalproex. Generalized seizures: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: carbamazepine, phenytoin, topiramate, levetiracetam, gabapentin, phenytoin, topiramate, levetiracetam, primidone or lamotrigine.

Prior Authorization Group Description:

PIQRAY

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Hormone receptor (HR)-positive (i.e., estrogen or progesterone receptor [ER/PR]-positive), HER2-negative, advanced (locally recurrent) or metastatic, and positive for PIK3CA-mutation.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

Prescribed in combination with fulvestrant after disease progression on or after an endocrine therapy (e.g., anastrozole, exemestane, fulvestrant, toremifene, letrozole, tamoxifen, or megestrol acetate).

Prior Authorization Group Description:

PLEGRIDY

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

PRALUENT

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

GENETICALLY MEDIATED PRIMARY HYPERLIPIDEMIA (INCLUDING HETEROZYGOUS FAMILIAL HYPERCHOLESTEROLEMIA): Confirmation (e.g., medical records, chart notes, laboratory values) of LDL level suggestive of a diagnosis of familial hypercholesterolemia (e.g., adults: LDL of 190 mg/dL or greater). NON-GENETICALLY MEDIATED PRIMARY HYPERLIPIDEMIA: Request meets both of the following (a and b): a) Confirmation of a LDL of 100 mg/dL or greater AND b) a diagnosis of secondary hyperlipidemia has been ruled out with confirmation of absence of all of the following potential causes of elevated cholesterol (a-e): a) hypothyroidism, b) obstructive liver disease, c) renal disease, d) nephrosis, e) medications which can increase lipid levels including, but not limited to: glucocorticoids, sex hormones, antipsychotics, antiretrovirals, immunosuppressive agents, retinoic acid derivatives. HYPERCHOLESTEROLEMIA WITH HISTORY OF CLINICAL ATHEROSCLEROTIC CARDIOVASCULAR DISEASE (ASCVD): Confirmation of an LDL of 70 mg/dL or greater AND history of clinical ASCVD defined as one of the following: Acute coronary syndromes, Myocardial infarction, Stable or unstable angina, Coronary or other arterial revascularization (e.g., percutaneous coronary intervention or coronary artery bypass graft surgery), Stroke, Peripheral artery disease presumed to be of atherosclerotic origin, Transient ischemic attack (TIA), Clinically significant coronary heart disease (CHD) diagnosed by invasive or noninvasive testing (such as coronary angiography, stress test using treadmill, stress echocardiography, or nuclear imaging), Carotid artery occlusion greater than 50% without symptoms, Renal artery stenosis or renal artery stent procedure. CONTINUATION OF THERAPY: Confirmation of LDL reduction while on Praluent therapy AND, if tolerated, confirmation of continued statin therapy at the maximally tolerated dose.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist.

Coverage Duration:

6 months.

Other Criteria:

Failure of two of the following at maximally tolerated doses, unless contraindicated or clinically significant adverse effects are experienced: atorvastatin, rosuvastatin, simvastatin, ezetimibe/simvastatin, pitavastatin, pravastatin, fluvastatin, or lovastatin.

Prior Authorization Group Description:

PRETOMANID

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Confirmed resistance to fluoroquinolones, unless contraindicated or clinically significant adverse effects are experienced. CONTINUATION OF THERAPY: Confirmation of delayed culture conversion and total duration of pretomanid therapy has not exceeded 9 months.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an expert in the treatment of tuberculosis.

Coverage Duration:

Initial: 6 months. Reauthorization: 3 months.

Other Criteria:

Prescribed in combination with bedaquiline and linezolid. CONTINUATION OF THERAPY: Member meets one of the following (a or b): a) Prescribed in combination with bedaquiline and linezolid OR b) If member has completed at least 4 weeks of linezolid therapy, member continues to receive pretomanid in combination with bedaquiline.

Prior Authorization Group Description:

PREVYMIS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Member is receiving pimozide or ergot alkaloids. Member is receiving cyclosporine co-administered with pitavastatin or simvastatin.

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncology, hematology, infectious disease, or transplant specialist.

Coverage Duration:

Through day 100 post-transplantation.

Other Criteria:

Failure of generic valacyclovir or generic ganciclovir, unless contraindicated, clinically significant adverse effects are experienced, or member is at high risk for CMV.

Prior Authorization Group Description:

PROCARDIA CAPSULES

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

CHRONIC STABLE ANGINA: Failure of ONE of the following, unless contraindicated or clinically significant adverse effects are experienced: nifedipine SR, amlodipine or nicardipine. VASOSPASTIC ANGINA: Failure of ONE of the following, unless contraindicated or clinically significant adverse effects are experienced: nifedipine SR or amlodipine.

Prior Authorization Group Description:

PROLASTIN C

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a pulmonologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

PROLIA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

PROSTATE CANCER: Receiving or has received androgen deprivation therapy [e.g., leuprolide (Lupron), bicalutamide (Casodex) or Nilandron]. BREAST CANCER: Receiving or has received adjuvant endocrine therapy [e.g., tamoxifen or aromatase inhibitors such as anastrozole (Arimidex), exemestane (Aromasin) or letrozole (Femara)].

Prior Authorization Group Description:

PROMACTA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Thrombocytopenia in Chronic Hepatitis C: Confirmation of current or planned interferon-based treatment of chronic hepatitis C.

Age Restrictions:

Prescriber Restrictions:

CHRONIC IMMUNE THROMBOCYTOPENIA, SEVERE APLASTIC ANEMIA: Prescribed by or in consultation with a hematologist. THROMBOCYTOPENIA IN CHRONIC HEPATITIS C: Prescribed by or in consultation with a hematologist, gastroenterologist, or an infectious disease specialist.

Coverage Duration:

12 months.

Other Criteria:

Chronic Immune (Idiopathic) Thrombocytopenia: Failure of a corticosteroid (e.g., prednisone, methylprednisolone, or dexamethasone), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

PROTOPIC

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Tacrolimus 0.1%: 16 years and older.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of two medium to high potency topical corticosteroids (e.g., amcinonide, fluticasone propionate, triamcinolone acetonide, betamethasone valerate, fluocinolone acetonide, hydrocortisone butyrate, mometasone furoate, desoximetasone, fluocinonide or betamethasone dipropionate), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

PROVIGIL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Multiple sclerosis-related fatigue.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

PURIXAN

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with oncologist or hematologist.

Coverage Duration:

12 months.

Other Criteria:

One of the following: Failure of mercaptopurine tablets, unless contraindicated or clinically significant adverse effects are experienced OR member has a swallowing disorder or an inability to swallow tablets or capsules..

Prior Authorization Group Description:

QINLOCK

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

GASTROINTESTINAL STROMAL TUMOR: For members with PDGFRA exon 18 mutation, failure of Ayvakit, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

QUALAQUIN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Babesiosis. Plasmodium vivax malaria.

Exclusion Criteria:

For the treatment or prevention of nocturnal leg cramps.

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Malaria: 7 days. Babesiosis: 7-10 days.

Other Criteria:

Plasmodium vivax malaria: Infection is chloroquine-resistant.

Prior Authorization Group Description:

RADICAVA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Forced vital capacity greater than or equal to 80%, disease duration of less than or equal to 2 years, functionally retains most activities of daily living (defined as a baseline revised ALS Functional Rating Scale (ALSFRS-R) score with greater than or equal to 2 points in each of the 12 items, meets diagnostic criteria of definite or probable amyotrophic lateral sclerosis (ALS) based on El Escorial revised criteria. CONTINUATION OF THERAPY: Member continues to retain most activities of daily living, forced vital capacity greater than or equal to 80%, and ALSFRS-R score with greater than or equal to 2 points in each of the 12 items.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

6 months.

Other Criteria:

Prescribed in combination with riluzole unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

RAYALDEE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Patient has stage 3 or 4 chronic kidney disease (CKD) and serum total 25-hydroxyvitamin D level less than 30 ng/mL.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

REBLOZYL

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

TRANSFUSION DEPENDENT BETA-THALASSEMIA: Total volume of transfusions exceeds 6 red blood cell units within the last 6 months. No transfusion free period for greater than or equal to 35 days within the last 6 months. CONTINUATION OF THERAPY, TRANSFUSION DEPENDENT BETA THALASSEMIA: Member meets one of the following (a or b): a) Member is responding positively to therapy as evidenced by at least a 33% reduction in transfusion burden from baseline, b) Request is for a dose increase. MYELODYSPLASTIC SYNDROMES (MDS): Member requires 2 or more RBC units per 8 weeks confirmed for at least the last 16 weeks. Member has either a ring sideroblast of at least 15% of erythroid precursors in bone marrow or ring sideroblast of at least 5% if SF3B1 mutation is present. Member does not have del(5q) cytogenetic abnormality. CONTINUATION OF THERAPY, MDS: Member meets one of the following (a or b): a) Member is responding positively to therapy as evidenced by a decreased transfusion burden, b) Request is for a dose increase.

Age Restrictions:

Prescriber Restrictions:

TRANSFUSION DEPENDENT BETA-THALASSEMIA: Prescribed by or in consultation with a hematologist. MDS: Prescribed by or in consultation with a hematologist or oncologist.

Coverage Duration:

TRANSFUSION DEPENDENT BETA-THALASSEMIA, MDS: Initial: 2 months. Reauthorization: 6 months.

Other Criteria:

MDS: Failure of an erythropoiesis-stimulating agent used in combination with a granulocyte colony stimulating factor, unless clinically significant adverse effects are experienced, all are contraindicated, or confirmation of current serum erythropoietin greater than 500 mU/mL.

Prior Authorization Group Description:

RELISTOR

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of Amitiza and Movantik, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

REMICADE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Wegener's Granulomatosis.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Psoriatic Arthritis/Plaque Psoriasis: Prescribed by or in consultation with a rheumatologist or dermatologist. Crohn's Disease/Ulcerative Colitis: Prescribed by or in consultation with a gastroenterologist. Rheumatoid Arthritis/Ankylosing Spondylitis: Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

Rheumatoid Arthritis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. Plaque Psoriasis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin.

Prior Authorization Group Description:

RENFLEXIS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Psoriatic Arthritis/Plaque Psoriasis: Prescribed by or in consultation with a rheumatologist or dermatologist. Crohn's Disease/Ulcerative Colitis: Prescribed by or in consultation with a gastroenterologist. Rheumatoid Arthritis/Ankylosing Spondylitis: Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

Rheumatoid Arthritis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. Plaque Psoriasis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin.

Prior Authorization Group Description:

REPATHA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

GENETICALLY MEDIATED PRIMARY HYPERLIPIDEMIA (INCLUDING HETEROZYGOUS OR HOMOZYGOUS FAMILIAL HYPERCHOLESTEROLEMIA): Confirmation (e.g., medical records, chart notes, laboratory values) of LDL level suggestive of a diagnosis of familial hypercholesterolemia (e.g., adults: LDL 190 mg/dL or greater). NON-GENETICALLY MEDIATED PRIMARY HYPERLIPIDEMIA: Request meets both of the following (a and b): a) Confirmation of a LDL of 100 mg/dL or greater AND b) a diagnosis of secondary hyperlipidemia has been ruled out with confirmation of absence of all of the following potential causes of elevated cholesterol (a-e): a) hypothyroidism, b) obstructive liver disease, c) renal disease, d) nephrosis, e) medications which can increase lipid levels including, but not limited to: glucocorticoids, sex hormones, antipsychotics, antiretrovirals, immunosuppressive agents, retinoic acid derivatives. HYPERCHOLESTEROLEMIA WITH HISTORY OF CLINICAL ATHEROSCLEROTIC CARDIOVASCULAR DISEASE (ASCVD): Confirmation of an LDL of 70 mg/dL or greater AND history of clinical ASCVD defined as one of the following: Acute coronary syndromes, Myocardial infarction, Stable or unstable angina, Coronary or other arterial revascularization (e.g., percutaneous coronary intervention or coronary artery bypass graft surgery), Stroke, Peripheral artery disease presumed to be of atherosclerotic origin, Transient ischemic attack (TIA), Clinically significant coronary heart disease (CHD) diagnosed by invasive or noninvasive testing (such as coronary angiography, stress test using treadmill, stress echocardiography, or nuclear imaging), Carotid artery occlusion greater than 50% without symptoms, Renal artery stenosis or renal artery stent procedure. CONTINUATION OF THERAPY: Confirmation of LDL reduction while on Repatha therapy AND, if tolerated, confirmation of continued statin therapy at the maximally tolerated dose.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a cardiologist, endocrinologist, or lipid specialist.

Coverage Duration:

6 months.

Other Criteria:

Failure of two of the following at maximally tolerated doses, unless contraindicated or clinically significant adverse effects are experienced: atorvastatin, rosuvastatin, simvastatin, ezetimibe/simvastatin, pitavastatin, pravastatin, fluvastatin, or lovastatin.

Prior Authorization Group Description:

RETEVMO

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

REVATIO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Members on concomitant nitrates (e.g., Nitrodur, Nitrobid, Nitrostat, Isordil, Ismo). Members on concomitant guanylate cyclase stimulator, such as riociguat (Adempas).

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

REVCOVI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an immunologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

REVLIMID

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

REXULTI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of aripiprazole and one of the following generic atypical antipsychotics, unless contraindicated or clinically significant adverse effects are experienced: risperidone, olanzapine, quetiapine, ziprasidone.

Prior Authorization Group Description:

RINVOQ

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin.

Prior Authorization Group Description:

RITUXIMAB

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

All oncology indications: Prescribed by or in consultation with an oncologist or hematologist. Rheumatoid arthritis, granulomatosis with polyangiitis, microscopic polyangiitis: Prescribed by or in consultation with a rheumatologist. Pemphigus vulgaris: Prescribed by or in consultation with a dermatologist.

Coverage Duration:

12 months.

Other Criteria:

Rheumatoid Arthritis: Prescribed in combination with methotrexate, unless contraindicated or clinically significant adverse effects were experienced with prior methotrexate therapy AND failure of Enbrel or Humira, unless contraindicated or clinically significant adverse effects are experienced. Granulomatosis with polyangiitis, Microscopic polyangiitis: Prescribed in combination with a glucocorticoid (e.g. prednisone, prednisolone, dexamethasone).

Prior Authorization Group Description:

ROZLYTREK

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

ROS1-POSITIVE NON-SMALL CELL LUNG CANCER: Confirmation of a ROS1 mutation. Member has not received prior ROS1 targeted therapy (e.g., Xalkori, Zykadia, Lorbrena). NTRK FUSION-POSITIVE SOLID TUMOR: Confirmation of an NTRK gene fusion (e.g., ETV6-NTRK3, TPM3-NTRK1). Member has not received prior NTRK targeted therapy (e.g., Vitrakvi).

Age Restrictions:

NTRK FUSION-POSITIVE SOLID TUMOR: Age greater than or equal to 12 years.

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

RUBRACA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

OVARIAN CANCER: Mutations in the BRCA genes OR member has a complete or partial response to two or more platinum-based chemotherapy regimens.

Age Restrictions:

Prescriber Restrictions:

PROSTATE CANCER: Prescribed by or in consultation with an oncologist or urologist. ALL OTHER ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

RUZURGI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Confirmation of a baseline clinical muscle strength assessment (examples may include but are not limited to the Quantitative Myasthenia Gravis (QMG) score, triple-timed up-and-go test (3TUG), Timed 25-foot Walk test (T25FW)). CONTINUATION OF THERAPY: Member is responding positively to therapy as evidenced by clinical muscle strength assessments.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

RYDAPT

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Acute Myeloid Leukemia: Positive for the FLT3 mutation.

Age Restrictions:

Prescriber Restrictions:

Acute Myeloid Leukemia: Prescribed by or in consultation with an oncologist or hematologist. Advanced Systemic Mastocytosis: Prescribed by or in consultation with an oncologist, allergist, or immunologist.

Coverage Duration:

12 months.

Other Criteria:

Acute Myeloid Leukemia: for induction therapy, prescribed in combination with cytarabine and daunorubicin OR for consolidation therapy, prescribed in combination with cytarabine.

Prior Authorization Group Description:

SAVELLA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Depression.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Fibromyalgia: Failure of duloxetine or Lyrica, unless contraindicated or clinically significant adverse effects are experienced. Depression: Failure of ONE of the following generic antidepressants, unless contraindicated or clinically significant adverse effects are experienced: bupropion, bupropion SR, bupropion XL, citalopram, desvenlafaxine succinate, duloxetine, escitalopram, fluoxetine, mirtazapine, sertraline, venlafaxine, or venlafaxine XR.

Prior Authorization Group Description:

SECUADO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Medical justification supports inability to use Saphris (asenapine sublingual tablets) (e.g., contraindications to excipients).

Prior Authorization Group Description:

SEROQUEL XR

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Schizophrenia: Failure of two of the following generic atypical antipsychotics, unless contraindicated or clinically significant adverse effects are experienced: risperidone, olanzapine, quetiapine immediate release, ziprasidone, aripiprazole.

Prior Authorization Group Description:

SILIQ

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a rheumatologist or dermatologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin.

Prior Authorization Group Description:

SIMPONI(prefilled syringe, auto-injector)

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

PSORIATIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist or dermatologist. ULCERATIVE COLITIS: Prescribed by or in consultation with a gastroenterologist. RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS: Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin.

Prior Authorization Group Description:

SIMPONI ARIA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

PSORIATIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist or dermatologist. RHEUMATOID ARTHRITIS, ANKYLOSING SPONDYLITIS: Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin.

Prior Authorization Group Description:

SKYRIZI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a dermatologist or rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine, or acitretin.

Prior Authorization Group Description:

SOMA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

SOMAVERT

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an endocrinologist.

Coverage Duration:

12 months.

Other Criteria:

Inadequate response to surgery or radiation therapy, unless surgery or radiation therapy is not appropriate for the patient.

Prior Authorization Group Description:

SOVALDI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Criteria will be applied consistent with current AASLD-IDSA guidance.

Exclusion Criteria:

Required Medical Information:

Refer to the document, "Recommendations for Testing, Managing, and Treating Hepatitis C," by AASLD-IDSA available at http://www.hcvguidelines.org for drug regimen and duration of treatment based on genotype, treatment status, previous drug regimens used, past medical history and comorbidities.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a gastroenterologist, hepatologist or infectious disease physician.

Coverage Duration:

Criteria will be applied consistent with current AASLD-IDSA guidance.

Other Criteria:

Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: Mavyret, Harvoni, Epclusa, Vosevi, and Zepatier for applicable genotypes.

Prior Authorization Group Description:

SPRAVATO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Currently on an oral antidepressant (must not be an agent previously tried and failed). CONTINUATION OF THERAPY: Member is responding positively to therapy and is using Spravato in combination with an oral antidepressant.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

Initial: 4 weeks. Reauthorization: 6 months.

Other Criteria:

Failure of two antidepressants (e.g., selective serotonin reuptake inhibitor [SSRI], serotonin-norepinephrine reuptake inhibitor [SNRI], tricyclic antidepressant [TCA], bupropion, mirtazapine) from two different classes, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

SPRITAM

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Medical justification must be provided why patient cannot take generic levetiracetam tablets or liquid.

Prior Authorization Group Description:

SPRYCEL

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

CHRONIC MYELOGENOUS LEUKEMIA, ACUTE LYMPHOBLASTIC LEUKEMIA: Confirmation that the member has Philadelphia chromosome positive disease.

Age Restrictions:

Prescriber Restrictions:

CHRONIC MYELOGENOUS LEUKEMIA, ACUTE LYMPHOBLASTIC LEUKEMIA: Prescribed by or in consultation with an oncologist or hematologist. ALL OTHER COVERED ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

GASTROINTESTINAL STROMAL TUMOR: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: imatinib, Sutent or Stivarga.

Prior Authorization Group Description:

STELARA IV

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a gastroenterologist.

Coverage Duration:

4 weeks.

Other Criteria:

CROHN'S DISEASE: Failure of Humira or Remicade and one of the following, unless contraindicated or clinically significant adverse effects are experienced: 6-mercaptopurine, azathioprine or methotrexate.

Prior Authorization Group Description:

STELARA SC

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

PSORIATIC ARTHRITIS, PLAQUE PSORIASIS: Prescribed by or in consultation with a rheumatologist or dermatologist. CROHN'S DISEASE, ULCERATIVE COLITIS: Prescribed by or in consultation with a gastroenterologist.

Coverage Duration:

12 months.

Other Criteria:

PLAQUE PSORIASIS: Failure of ONE of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine, or acitretin. PSORIATIC ARTHRITIS: Failure of one TNF inhibitor (e.g., Humira, Enbrel, Cimzia, Simponi, Simponi Aria, Remicade, Inflectra), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

STIVARGA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

STRENSIQ

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

SUBSYS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Age 18 or greater.

Prescriber Restrictions:

Coverage Duration:

Through the end of the Plan contract year.

Other Criteria:

Patient is already taking and is tolerant to around-the-clock opioid therapy. Patients are considered opioid tolerant when taking another opioid daily for a week or longer (for example, at least 60 mg of oral morphine per day or an equianalgesic dose of another opioid).

Prior Authorization Group Description:

SUNOSI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

NARCOLEPSY: Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of armodafinil (Nuvigil) or modafinil (Provigil), unless contraindicated or clinically significant side effects are experienced.

Prior Authorization Group Description:

SURMONTIL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Irritable bowel syndrome.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Depression: Failure of one of the following generic antidepressants, unless contraindicated or clinically significant adverse effects are experienced: bupropion, bupropion SR, bupropion XL, citalopram, desvenlafaxine succinate, duloxetine, escitalopram, fluoxetine, mirtazapine, sertraline, venlafaxine, or venlafaxine XR.

Prior Authorization Group Description:

SYMDEKO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Presence of homozygous F508del mutation or at least one mutation in the CFTR gene that is responsive to tezacaftor/ivacaftor.

Age Restrictions:

Age greater than or equal to 6 years.

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

SYMLINPEN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Previous use of mealtime insulin therapy or an insulin pump.

Prior Authorization Group Description:

SYMPAZAN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Other Criteria:

Medical justification supports inability to use clobazam tablets and oral suspension (e.g., contraindications to excipients).

Prior Authorization Group Description:

TABRECTA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

NON-SMALL CELL LUNG CANCER: Disease is epidermal growth factor receptor (EGFR) wild-type and anaplastic lymphoma kinase (ALK) negative.

Age Restrictions:

Prescriber Restrictions:

ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

TAFAMIDIS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

TRANSTHYRETIN AMYLOID CARDIOMYOPATHY (ATTR-CM): Member has not had a liver transplant. Diagnosis is supported by (a or b): a) Confirmation of amyloid deposition on biopsy and either transthyretin (TTR) precursor protein (e.g., by immunohistochemistry, scintigraphy, mass spectrometry) or a TTR mutation by genetic testing. b) Member meets all of the following (i, ii, and iii): i) Echo, CMR, or PET findings are consistent with cardiac amyloidosis, AND ii) Cardiac update is Grade 2 or 3 on a radionuclide scan utilizing one of the following radiotracers (1, 2, or 3): 1) 99m technetium (Tc)-labeled 3,3-diphosphono-1,2-propanodicarboxylic acid (DPD), 2) 99mTc-labeled pyrophosphate (PYP), or 3) 99mTc-labeled hydroxymethylene diphosphonate (HMDP), AND iii) Each of the following laboratory tests is negative for monoclonal protein (1, 2, and 3): 1) Serum kappa/lambda free light chain ratio analysis, 2) Serum protein immunofixation, 3) Urine protein immunofixation. CONTINUATION OF THERAPY, ATTR-CM: Maintained on therapy with positive response, including but not limited to, improvement or stabilization in any of the following parameters: 1) walking ability, 2) nutrition (e.g., body mass index), 3) cardiac related hospitalization, 4) cardiac procedures or laboratory tests (e.g., Holter monitoring, echocardiography, electrocardiogram, plasma BNP or NT-proBNP, serum troponin).

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a cardiologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

TAGRISSO

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

NON-SMALL CELL LUNG CANCER: Disease is positive for either of the following (a or b): a) sensitizing EGFR mutation (e.g., exon 19 deletion or insertion, exon 21 point mutation (L858R, L861Q), exon 18 point mutation (G719X), exon 20 point mutation (S768I)), OR b) T790M mutation.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

TAKHZYRO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Age greater than or equal to 12 years.

Prescriber Restrictions:

Prescribed by or in consultation with an immunologist, allergist, hematologist, or rheumatologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

TALTZ

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

PLAQUE PSORIASIS, PSORIATIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist or dermatologist. ANKYLOSING SPONDYLITIS: Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

PLAQUE PSORIASIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine or acitretin. PSORIATIC ARTHRITIS: Failure of one TNF inhibitor (e.g., Humira, Enbrel, Cimzia, Simponi, Simponi Aria, Remicade, Inflectra), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

TALZENNA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Confirmation of human epidermal growth factor receptor 2 (HER2)-negative disease and mutation in the BRCA genes.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

TARCEVA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

NON-SMALL CELL LUNG CANCER: Disease is positive for a sensitizing EGFR mutation (e.g., exon 19 deletion or insertion, exon 21 point mutation (L858R, L861Q), exon 18 point mutation (G719X), exon 20 point mutation (S768I)). RENAL CELL CARCINOMA: Confirmation of non-clear cell histology.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

PANCREATIC CANCER: Prescribed in combination with gemcitabine.

Prior Authorization Group Description:

TARGRETIN GEL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D. Some medically-accepted indications.

Off Label Uses:

Primary cutaneous B-cell lymphoma types: primary cutaneous marginal zone lymphoma and primary cutaneous follicle center lymphoma.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

TASIGNA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

CHRONIC MYELOGENOUS LEUKEMIA, ACUTE LYMPHOBLASTIC LEUKEMIA: Confirmation that the member has Philadelphia chromosome positive disease.

Age Restrictions:

Prescriber Restrictions:

CHRONIC MYELOGENOUS LEUKEMIA, ACUTE LYMPHOBLASTIC LEUKEMIA: Prescribed by or in consultation with an oncologist or hematologist. GASTROINTESTINAL STROMAL TUMOR: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

GASTROINTESTINAL STROMAL TUMOR: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: imatinib, Sutent or Stivarga.

Prior Authorization Group Description:

TAVALISSE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of a corticosteroid (e.g., prednisone, methylprednisolone, or dexamethasone), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

TAZVERIK

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

EPITHELIOID SARCOMA: Tumor demonstrates loss of INI1 expression through inactivation, deletion, or mutation of the INI1 (SMARCB-1) gene.

Age Restrictions:

EPITHELIOID SARCOMA: Age 16 years or older.

Prescriber Restrictions:

FOLLICULAR LYMPHOMA: Prescribed by or in consultation with an oncologist or hematologist. EPITHELIOID SARCOMA: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

TECENTRIQ

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

EXTENSIVE-STAGE SMALL CELL LUNG CANCER: Prescribed in combination with carboplatin and etoposide. TRIPLE NEGATIVE BREAST CANCER: Hormone-receptor (HR)-negative, estrogen-receptor (ER)-negative, and human epidermal growth factor receptor 2 (HER2)-negative disease. Prescribed in combination with protein-bound paclitaxel (nab-paclitaxel). Tumor expresses PD-L1.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

NON-SMALL CELL LUNG CANCER: If a known epidermal growth factor receptor (EGFR) or anaplastic lymphoma kinase (ALK) genomic tumor aberration exists, then for ALK+ disease: prior trial of Xalkori, Alecensa, or Zykadia OR for EGFR+ disease: prior trial of Tarceva, Gilotrif, or Iressa.

Prior Authorization Group Description:

TECFIDERA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

TEGSEDI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Confirmation of transthyretin (TTR) mutation. Confirmation of amyloid deposition on biopsy or medical justification is provided as to why treatment should be initiated in the presence of a negative biopsy or no biopsy. Member has not had a liver transplant. CONTINUATION OF THERAPY: Maintained on therapy with positive response, including but not limited to improvement in any of the following parameters: 1) neuropathy (motor function, sensation, reflexes, walking ability), 2) nutrition (body mass index), 3) cardiac parameters (Holter monitoring, echocardiography, electrocardiogram, plasma BNP or NT-proBNP, serum troponin), 4) renal parameters (creatinine clearance, urine albumin), 5) ophthalmic parameters (eye exam).

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

TENEX

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: amlodipine/benazepril, benazepril/hydrochlorothiazide, captopril, captopril/hydrochlorothiazide, fosinopril, fosinopril/hydrochlorothiazide, lisinopril, lisinopril/hydrochlorothiazide, quinapril, quinapril/hydrochlorothiazide, losartan, losartan/hydrochlorothiazide, valsartan, valsartan/hydrochlorothiazide, irbesartan, irbesartan/hydrochlorothiazide, candesartan, candesartan/hydrochlorothiazide, carvedilol, labetalol, acebutolol, atenolol, bisoprolol, bisoprolol/hydrochlorothiazide, timolol, nadolol, propranolol, metoprolol, metoprolol, nifedipine SR, amlodipine, nicardipine.

Prior Authorization Group Description:

TEPEZZA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

THYROID EYE DISEASE (TED): Member has active TED with a clinical activity score (CAS) of at least 4 or greater. Member is euthyroid with confirmation of a recent (within the last 30 days) free thyroxine (FT4) and free triiodothyronine (FT3) levels within the laboratory defined reference range. Member has not had previous surgical intervention for TED. Member does not require surgical ophthalmological intervention. Member has not received 8 or more infusions lifetime (including the initial 10 mg/kg first infusion). CONTINUATION OF THERAPY, TED: Member is responding positively to therapy as evidenced by both of the following (a and b): a) at least a 2 mm or greater reduction in proptosis AND b) at least a 2 point or greater reduction in CAS from baseline. Member does not require surgical ophthalmological intervention. Iffetime (including the initial 10 mg/kg first infusion).

Age Restrictions:

Prescriber Restrictions:

TED: Prescribed by or in consultation with an ophthalmologist.

Coverage Duration:

6 months.

Other Criteria:

TED: Failure of a systemic corticosteroid, unless clinically significant adverse effects are experienced or all are contraindicated.

Prior Authorization Group Description:

TETRABENAZINE

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

TIBSOVO

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Presence of an isocitrate dehydrogenase-1 (IDH1) mutation. For newly diagnosed acute myeloid leukemia (AML), member is age 60 years or older OR medical justification supports inability to use intensive induction therapy.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Other Criteria:

For age less than 60 years where medical justification does not support inability to use intensive induction therapy, disease has relapsed or is refractory following treatment with standard antineoplastic induction agents (e.g., cytarabine, idarubicin, daunorubicin, cladribine, midostaurin).

Prior Authorization Group Description:

TOLSURA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Hematologic malignancy for prophylaxis of aspergillosis or candidiasis.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

HISTOPLASMOSIS: 6 weeks. ASPERGILLOSIS: 3 months. BLASTOMYCOSIS, HEMATOLOGIC MALIGNANCY: 6 months.

Other Criteria:

ALL INDICATIONS: Failure of generic itraconazole capsule, unless contraindicated or clinically significant adverse effects are experienced. ASPERGILLOSIS: Failure of voriconazole, unless contraindicated or clinically significant adverse effects are experienced. HEMATOLOGIC MALIGNANCY: For candidiasis prophylaxis, failure of fluconazole, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

TREMFYA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a rheumatologist or dermatologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of ONE of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, cyclosporine, or acitretin.

Prior Authorization Group Description:

TRIHEXYPHENIDYL

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prior authorization is required for patients 65 years and older. Prior authorization is not required for patients 64 years and younger.

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Parkinsons disease/Parkinsonism: Failure of two of the following, unless contraindicated or clinically significant adverse effects are experienced: amantadine, levodopa/carbidopa, entacapone, pramipexole, ropinirole, selegiline.

Prior Authorization Group Description:

TRIKAFTA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

CYSTIC FIBROSIS: Diagnosis of cystic fibrosis (CF) confirmed by both of the following (a and b): a) Clinical symptoms consistent with CF in at least one organ system, or positive newborn screen or genetic testing for siblings of patients with CF, AND b) Evidence of cystic fibrosis transmembrane conductance regulator (CFTR) dysfunction confirmed by one of the following (i or ii): i) Evidence of clinical severity as defined by an average sweat chloride greater than 60 mmol/L, OR ii) Genetic testing confirming the presence of two disease-causing mutations in CFTR gene, one from each parent allele, and one of which is a F508del mutation. Confirmation that pulmonary function tests, performed within the last 90 days, show a percent predicted forced expiratory volume in 1 second (ppFEV1) that is between 40-90%. Trikafta is not prescribed concurrently with other CFTR modulators (e.g., Orkambi, Kalydeco, Symdeko). CONTINUATION OF THERAPY, CYSTIC FIBROSIS: For members that received at least 12 weeks of therapy, member is responding positively to therapy as evidenced by stabilization in ppFEV1 if baseline was 70% or greater or increase in ppFEV1 if baseline was less than 70%.

Age Restrictions:

CYSTIC FIBROSIS: Age greater than or equal to 12 years.

Prescriber Restrictions:

CYSTIC FIBROSIS: Prescribed by or in consultation with a pulmonologist.

Coverage Duration:

Initial: 4 months. Reauthorization: 12 months.

Prior Authorization Group Description:

TUKYSA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

ALL ONCOLOGY INDICATIONS: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

TURALIO

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

TYMLOS

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Total duration of therapy on parathyroid hormone (PTH) analogs (e.g., Tymlos, Forteo) has not exceeded 2 years.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Member meets one of the following (a, b, or c): a) Failure of a bisphosphonate (e.g., alendronate) unless contraindicated or clinically significant adverse effects are experienced. OR b) Bone mineral density T-score at hip or spine is -3.5 or less. OR c) Bone mineral density T-score at hip or spine is -2.5 or less with a history of major osteoporotic fracture of the hip, spine, forearm, wrist, or humerus.

Prior Authorization Group Description:

TYSABRI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

MULTIPLE SCLEROSIS: Prescribed by or in consultation with a neurologist. CROHN'S DISEASE: Prescribed by or in consultation with a gastroenterologist.

Coverage Duration:

12 months.

Other Criteria:

RELAPSING-REMITTING MULTIPLE SCLEROSIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Aubagio, Tecfidera, Gilenya, Avonex, Betaseron, Plegridy, glatiramer, Copaxone, Glatopa, Extavia or Rebif. CROHN'S DISEASE: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: Humira or Remicade.

Prior Authorization Group Description:

ULTRAVATE LOTION

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of generic halobetasol propionate and generic clobetasol propionate, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

UPTRAVI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

VALCHLOR

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following skin-directed therapies unless contraindicated or clinically significant adverse effects are experienced: topical corticosteroid (e.g., betamethasone, clobetasol), topical retinoid (e.g., Targretin, Avage, Fabior, Tazorac), topical imiquimod (Aldara).

Prior Authorization Group Description:

VALTOCO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Diagnosis of partial or generalized epilepsy.

Age Restrictions:

6 years of age or older.

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Other Criteria:

Medical justification supports inability to use diazepam rectal gel (e.g., contraindications to excipients).

Prior Authorization Group Description:

VANCOCIN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

C. Diff diarrhea: 14 days. Staph enterocolitis: 10 days. Recurrent C. Diff: 12 weeks.

Prior Authorization Group Description:

VENCLEXTA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

AML: Age 60 years or greater, OR medical justification supports inability to use intensive induction chemotherapy. Prescribed in combination with azacitidine, decitabine, or low-dose cytarabine.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Other Criteria:

MANTLE CELL LYMPHOMA: Failure of at least one previous therapy (e.g., a Rituxan based regimen), unless contraindicated or clinically significant adverse effects are experienced. CLL/SLL: Request meets one of the following (a or b): a) Prescribed in combination with Gazyva as first-line therapy OR b) Failure of at least one previous therapy (e.g., Imbruvica, Campath, or high-dose methylprednisolone with Rituxan), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

VERSACLOZ

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Psychotic disorder associated with Parkinson's disease.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of clozapine (Clozaril) or FazaClo, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

VERZENIO

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative advanced or metastatic breast cancer.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

Prescribed as a single agent or in combination with an aromatase inhibitor (e.g., letrozole, anastrozole or exemestane) or fulvestrant. For men receiving an aromatase inhibitor: Prescribed in combination with an agent that suppresses testicular steroidogenesis (e.g., gonadotropin-releasing hormone agonists).

Prior Authorization Group Description:

VIBERZI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of loperamide unless contraindicated or clinically significant adverse effects are experienced AND For members 64 years and younger, failure of diphenoxylate-atropine (Lomotil) or dicyclomine, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

VIMOVO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: pantoprazole, lansoprazole or omeprazole AND For osteoarthritis or rheumatoid arthritis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: ibuprofen, diclofenac sodium or potassium, etodolac, fenoprofen, ketoprofen, meloxicam, naproxen, oxaprozin, piroxicam, salsalate, sulindac, tolmetin OR For ankylosing spondylitis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: diclofenac sodium, naproxen or sulindac OR For juvenile idiopathic arthritis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: diclofenac sodium, naproxen or sulindac OR For juvenile idiopathic arthritis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: diclofenac sodium, naproxen or sulindac OR For juvenile idiopathic arthritis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: diclofenac sodium, naproxen or sulindac OR For juvenile idiopathic arthritis: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: etodolac, ibuprofen, meloxicam, naproxen, oxaprozin, tolmetin.

Prior Authorization Group Description:

VINBLASTINE

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Confirmation that vinblastine is being used as palliative therapy.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

VINCRISTINE

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Patients with the demyelinating form of Charcot-Marie-Tooth syndrome.

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

VITRAKVI

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Known acquired tropomyosin receptor kinase resistance mutation.

Required Medical Information:

Confirmation of positive neurotrophic receptor tyrosine kinase gene fusion mutation.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

VIZIMPRO

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Disease is positive for a sensitizing EGFR mutation (e.g., exon 19 deletion or insertion, exon 21 point mutation (L858R, L861Q), exon 18 point mutation (G719X), exon 20 point mutation (S768I)).

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

VOSEVI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

If cirrhosis is present, confirmation of Child-Pugh A status. Refer to the document, "Recommendations for Testing, Managing, and Treating Hepatitis C," by AASLD-IDSA available at http://www.hcvguidelines.org for drug regimen and duration of treatment based on genotype, treatment status, previous drug regimens used, past medical history and comorbidities.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a gastroenterologist, hepatologist or infectious disease physician.

Coverage Duration:

12 weeks.

Other Criteria:

Criteria will be applied consistent with current AASLD-IDSA guidance.

Prior Authorization Group Description:

VOTRIENT

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

VRAYLAR

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of TWO of the following atypical antipsychotics, unless contraindicated or clinically significant adverse effects are experienced: aripiprazole, ziprasidone, quetiapine, olanzapine, risperidone.

Prior Authorization Group Description:

VUMERITY

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

MULTIPLE SCLEROSIS: Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

VYONDYS 53

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

DUCHENNE MUSCULAR DYSTROPHY (DMD): DMD with mutation amenable to exon 53 skipping confirmed by genetic testing.

Age Restrictions:

Prescriber Restrictions:

DMD: Prescribed by or in consultation with a neurologist.

Coverage Duration:

6 months.

Other Criteria:

DMD: Currently stable on an oral corticosteroid regimen (e.g., prednisone), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

WAKIX

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

NARCOLEPSY: Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Other Criteria:

NARCOLEPSY: Failure of Sunosi, unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

XALKORI

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

NON-SMALL CELL LUNG CANCER: Disease is ALK, ROS1, or MET positive. INFLAMMATORY MYOFIBROBLASTIC TUMOR, ANAPLASTIC LARGE CELL LYMPHOMA: Disease is ALK-positive.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

XATMEP

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Less than 18 years of age.

Prescriber Restrictions:

ACUTE LYMPHOCYTIC LEUKEMIA: Prescribed by or in consultation with an oncologist or hematologist. POLYARTICULAR JUVENILE IDIOPATHIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist.

Coverage Duration:

12 months.

Other Criteria:

Medical justification as to why member cannot use methotrexate tablets.

Prior Authorization Group Description:

XCOPRI

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a neurologist.

Coverage Duration:

12 months.

Other Criteria:

Failure of two of the following generic antiepileptic drugs, unless contraindicated or clinically significant adverse effects are experienced: lamotrigine, topiramate, oxcarbazepine, carbamazepine, phenytoin, valproic acid, divalproex sodium, felbamate, gabapentin, levetiracetam, pregabalin, tiagabine, zonisamide.

Prior Authorization Group Description:

XELJANZ

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

RHEUMATOID ARTHRITIS: Prescribed by or in consultation with a rheumatologist. PSORIATIC ARTHRITIS: Prescribed by or in consultation with a rheumatologist or dermatologist. ULCERATIVE COLITIS (IMMEDIATE-RELEASE ONLY): Prescribed by or in consultation with a gastroenterologist.

Coverage Duration:

12 months.

Other Criteria:

RHEUMATOID ARTHRITIS: Failure of one of the following, unless contraindicated or clinically significant adverse effects are experienced: methotrexate, sulfasalazine, hydroxychloroquine, d-penicillamine, azathioprine or auranofin. PSORIATIC ARTHRITIS: Failure of one TNF inhibitor (e.g., Humira, Enbrel, Cimzia, Simponi, Simponi Aria, Remicade, Inflectra), unless predominantly axial disease, contraindicated, or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

XENLETA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Request is for continuation of therapy initiated in an acute care hospital from which member was discharged OR culture and sensitivity (C&S) report for the current infection shows isolated pathogen is an organism susceptible to Xenleta, unless provider confirms that obtaining a C&S report is not feasible.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

7 days.

Other Criteria:

For members initiating Xenleta therapy outside of an acute care hospital, one of the following (a, b, or c): a) If a C&S report is available: Failure of 2 antibiotics to which the isolated pathogen is susceptible (if available) per C&S report, unless all are contraindicated or clinically significant adverse effects are experienced. b) C&S report shows resistance or lack of susceptibility of the isolated pathogen to all formulary antibiotics FDA-approved for member's diagnosis. c) If provider confirms that obtaining a C&S report is not feasible: Failure of 2 antibiotics indicated for member's diagnosis (if available), unless all are contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

XEOMIN

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

XERMELO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Prescribed in combination with a somatostatin analog (e.g., octreotide, lanreotide) unless contraindicated or clinically significant adverse effects are experienced.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Other Criteria:

Failure of a somatostatin analog (e.g., octreotide, lanreotide) unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

XOLAIR

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

ASTHMA: Positive skin test or in vitro reactivity to a perennial aeroallergen AND immunoglobulin E (IgE) level greater than or equal to 30 IU/mL.

Age Restrictions:

ASTHMA: 6 years of age or older. CHRONIC IDIOPATHIC URTICARIA: 12 years of age or older.

Prescriber Restrictions:

ASTHMA: Prescribed by or in consultation with a pulmonologist, immunologist, or allergist. CHRONIC IDIOPATHIC URTICARIA: Prescribed by or in consultation with an allergist, dermatologist, or immunologist.

Coverage Duration:

12 months.

Other Criteria:

ASTHMA: Failure of one inhaled corticosteroid (e.g., beclomethasone, budesonide, flunisolide, fluticasone, mometasone, ciclesonide), unless contraindicated or clinically significant adverse effects are experienced. CHRONIC IDIOPATHIC URTICARIA: Failure of one H1 Antihistamine (e.g., levocetirizine or desloratadine), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

XOSPATA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Confirmation of the presence of a FLT3 mutation.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

XPOVIO

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

MULTIPLE MYELOMA: Member has received at least 4 prior lines of therapy that include all of the following (a, b, and c): a) Two proteasome inhibitors (e.g., bortezomib, Kyprolis, Ninlaro), b) Two immunomodulatory agents (e.g., Revlimid, pomalidomide, Thalomid), c) One anti-CD38 monoclonal antibody (e.g., Darzalex).

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or hematologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

XTANDI

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

PROSTATE CANCER: Concurrent use of a gonadotropin-releasing hormone (GnRH) analog or past bilateral orchiectomy.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or urologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

YERVOY

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

HEPATOCELLULAR CARCINOMA (new starts only): Member has not had previous treatment with a checkpoint inhibitor (e.g., Opdivo, Keytruda, Tecentriq, Imfinzi). NON-SMALL CELL LUNG CANCER (new starts only): Member has not previously progressed on a PD-1/PD-L1 inhibitor (e.g., Opdivo, Keytruda, Tecentriq, Imfinzi).

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

SMALL CELL LUNG CANCER, MALIGNANT PLEURAL MESOTHELIOMA: Failure of a platinum-containing regimen (e.g., cisplatin, carboplatin), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

YONSA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or urologist.

Coverage Duration:

12 months.

Other Criteria:

Medical justification supports inability to use Zytiga. Prescribed in combination with methylprednisolone. Member has previously had bilateral orchiectomy, failed androgen deprivation therapy (ADT) or will use ADT concurrently with Yonsa.

Prior Authorization Group Description:

ZALTRAP

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

Prescribed in combination with irinotecan or FOLFIRI (5-fluorouracil, leucovorin, and irinotecan). Previous treatment with one of the following: oxaliplatin-containing regimen (e.g., FOLFIRI, FOLFOX [leucovorin, 5-fluorouracil, oxaliplatin], CapeOX [capecitabine, oxaliplatin]) OR 5-fluorouracil and leucovorin containing regimen OR capecitabine containing regimen.

Prior Authorization Group Description:

ZARXIO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Myelodysplastic syndrome. Neutropenia in patients with HIV/AIDS. Hematopoietic syndrome of acute radiation syndrome.

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

12 months.

Prior Authorization Group Description:

ZEJULA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

OVARIAN, FALLOPIAN, OR PRIMARY PERITONEAL CANCER: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

ZELBORAF

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

MELANOMA, ERDHEIM-CHESTER DISEASE: Positive for the BRAF V600 mutation. NON-SMALL CELL LUNG CANCER, COLORECTAL CANCER: Positive for the BRAF V600E mutation. DIFFERENTIATED THYROID CARCINOMA: Positive for the BRAF mutation.

Age Restrictions:

Prescriber Restrictions:

ERDHEIM-CHESTER DISEASE, HAIRY CELL LEUKEMIA: Prescribed by or in consultation with an oncologist or hematologist. ALL OTHER ONCOLOGY INDICATION: Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Other Criteria:

NON-SMALL CELL LUNG CANCER: Failure of Tafinlar or Mekinist, unless contraindicated or clinically significant adverse effects are experienced. COLORECTAL CANCER: Failure of irinotecan or platinum-based therapy (e.g., oxaliplatin), unless contraindicated or clinically significant adverse effects are experienced.

Prior Authorization Group Description:

ZEPATIER

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

If cirrhosis is present, confirmation of Child-Pugh A status. For genotype 1a, confirmation of presence or absence of NS5A resistance-associated polymorphisms. Refer to the document, "Recommendations for Testing, Managing, and Treating Hepatitis C," by AASLD-IDSA available at http://www.hcvguidelines.org for drug regimen and duration of treatment based on genotype, treatment status, previous drug regimens used, past medical history and comorbidities.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a gastroenterologist, hepatologist or infectious disease physician.

Coverage Duration:

12 to 16 wks based on genotype, presence of NS5A resistance-associated polymorphisms, prior treatment.

Prior Authorization Group Description:

ZINPLAVA

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Confirmation of positive Clostridium difficile test.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

4 weeks.

Other Criteria:

Will receive or is currently receiving antibacterial drug treatment for Clostridium difficile infection (e.g., metronidazole, vancomycin, fidaxomicin) concomitantly with Zinplava. Has received appropriate treatment for past CDI recurrences, including a pulsed vancomycin regimen.

Prior Authorization Group Description:

ZULRESSO

Prior Authorization Indication:

All FDA-approved indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

No more than 6 months have passed since member has given birth.

Age Restrictions:

Prescriber Restrictions:

Coverage Duration:

4 weeks.

Other Criteria:

Failure of one of the following oral antidepressants, unless contraindicated or clinically significant adverse effects are experienced: selective serotonin reuptake inhibitor (SSRI), serotonin-norepinephrine reuptake inhibitor (SNRI), tricyclic antidepressant (TCA), bupropion, mirtazapine.

Prior Authorization Group Description:

ZYDELIG

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with a hematologist or oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

ZYKADIA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

NON-SMALL CELL LUNG CANCER: Disease is ALK or ROS1 positive. If disease is ROS1 positive, Zykadia is prescribed as first-line therapy. INFLAMMATORY MYOFIBROBLASTIC TUMOR: Disease is ALK-positive.

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist.

Coverage Duration:

12 months.

Prior Authorization Group Description:

ZYTIGA

Prior Authorization Indication:

All medically accepted indications not otherwise excluded from Part D.

Off Label Uses:

Exclusion Criteria:

Required Medical Information:

Age Restrictions:

Prescriber Restrictions:

Prescribed by or in consultation with an oncologist or urologist.

Coverage Duration:

12 months.

Other Criteria:

Prescribed in combination with prednisone. Member has previously had bilateral orchiectomy, failed androgen deprivation therapy (ADT) or will use ADT concurrently with abiraterone.

ABSTRAL	1
ACTEMRA IV	2
ACTEMRA SC	3
ACTHAR HP	4
ACTIQ	5
ACYCLOVIR	6
ADAKVEO	7
ADCIRCA	8
ADEMPAS	9
AFINITOR	10
AIMOVIG	
AJOVY	12
ALECENSA	
ALUNBRIG	14
AMANTADINE	
AMITRIPTYLINE	
AMITRIPTYLINE/CHLORDIAZEPOXIDE	17
AMITRIPTYLINE/PERPHENAZINE	18
AMPHOTERICIN B	
AMPYRA	20
ANTIHISTAMINES	
ANTIHISTAMINE COMBINATIONS	22
ARANESP	23
ARIKAYCE	24
AUBAGIO	
AUSTEDO	26
AVASTIN	
AYVAKIT	28
BALVERSA	29
BAXDELA	30
BELEODAQ	
BELSOMRA	32
BENLYSTA	
BENZTROPINE	
BEOVU	
BLEOMYCIN	
BOSULIF	
ВОТОХ	
BRAFTOVI	
BRIVIACT	
BRUKINSA	
C1 ESTERASE INHIBITOR	
CABLIVI	
САВОМЕТҮХ	
CALQUENCE	
САРLYTА	
CAPRELSA	47

CARISOPRODOL/ASPIRIN/CODEINE .	48
CAYSTON	49
CERDELGA	50
CEREZYME	51
CHLORZOXAZONE	52
CHORIONIC GONADOTROPIN	53
CIALIS	54
CIMZIA	55
CINQAIR	56
CLADRIBINE	57
CLOMIPRAMINE	58
COMETRIQ	59
COPIKTRA	60
COSENTYX	61
COTELLIC	62
CRINONE	63
CROFELEMER	64
CRYSVITA	65
CYCLOBENZAPRINE HCL	66
CYTARABINE	67
DAURISMO	68
DICLOFENAC GEL	69
DIPYRIDAMOLE	70
DISOPYRAMIDE	71
DOPTELET	72
DOXEPIN	73
DOXEPIN CREAM	74
DUEXIS	75
ELIDEL	76
EMEND 40 MG	77
EMFLAZA	78
EMGALITY	79
ENBREL	80
ENDARI	81
ENTRESTO	82
ENTYVIO	83
EPCLUSA	84
EPIDIOLEX	85
EPOETIN	86
ERGOLOID MESYLATES	87
ERLEADA	88
ESBRIET	89
ESTROGENS(Duavee, Divigel, Climara, , Mimvey, Prempro, Amabelz, Combipato Climara Pro, Evamist, Estrace, Activella, Menostar, Vivelle-Dot, Femhrt, Mimvey J Alora, Lopreeza, Premarin, Premphase).	h, Lo,
EVZIO	
EXONDYS 51	92

EYLEA	93	IN
FARYDAK	94	IN
FASENRA	95	JA
FENTORA	96	JU
FERRIPROX	97	JU
FINTEPLA	98	JY
FIORICET WITH CODEINE	99	KA
FIORINAL WITH CODEINE	100	KA
FIRAZYR	101	KA
FIRDAPSE	102	KA
FLECTOR	103	KE
FLUOROURACIL	104	KE
FORTEO	105	KE
GALAFOLD	106	KE
GANCICLOVIR		KE
GATTEX	108	KI
GAVRETO	109	KI
GILENYA	110	KC
GILOTRIF	111	KC
GLATIRAMER	112	KC
GLIMEPIRIDE		KU
GLYBURIDE	114	LA
GLYBURIDE/METFORMIN	115	LA
GRANIX	116	LE
HARVONI	117	LE
HERCEPTIN	118	LE
HETLIOZ	119	LII
HUMAN GROWTH HORMONE	120	LC
HUMIRA	121	LC
HYDROCODONE	122	LC
HYDROXYZINE HCL INJECTION	123	LU
HYDROXYZINE HCL ORAL		LY
HYDROXYZINE PAMOATE	125	MA
ICLUSIG	126	MA
IDHIFA	127	M
ILARIS	128	MI
ILUMYA	129	MI
IMATINIB	130	MI
IMBRUVICA	131	MI
IMIPRAMINE	132	MI
INDOMETHACIN	133	MI
INFLECTRA	134	MI
INGREZZA	135	MI
INLYTA	136	MI
INQOVI	137	MI
INREBIC	138	M
INTERFERON BETA-1A	139	M

INTERFERON BETA-1B140	
INTUNIV	
JAKAFI142	
JUBLIA143	
JUXTAPID144	
JYNARQUE145	
KADCYLA146	
KADIAN	
KALYDECO 148	
KAZANO149	
KERYDIN 150	
KETOROLAC TROMETHAMINE 151	
KEVEYIS152	
KEVZARA153	
KEYTRUDA 154	
KINERET155	
KISQALI(Kisqali Femara Co-Pack , Kisqali) 156	
KOMBIGLYZE XR 157	
KORLYM158	
KOSELUGO159	
KUVAN	
LATUDA161	
LAZANDA162	
LEMTRADA163	
LENVIMA164	
LEUKINE165	
LIDODERM166	
LONSURF	
LORBRENA168	
LOTRONEX	
LUCEMYRA	
LYNPARZA TABLET	
MAVENCLAD	
MAVYRET	
MAYZENT	
MEGACE	
MEGACE ES	
MEKINIST	
MEKTOVI	
MEPROBAMATE	
METAXALONE	
METHOCARBAMOL181	
METHOTREXATE INJ	
METHYLDOPA	
MIRVASO	
MOZOBIL	
MULPLETA	

NAMENDA	187]
NATPARA	188]
NAYZILAM	189]
NERLYNX	190]
NESINA	191]
NEULASTA	192]
NEUPOGEN	193	(
NINLARO	194	(
NIVESTYM	195	1
NORTHERA	196	1
NUBEQA	197]
NUCALA	198]
NUEDEXTA	199]
NUPLAZID	200	1
NUVIGIL	201	1
NUZYRA	202]
OCALIVA]
OCREVUS	204]
ODOMZO	. 205	
OFEV	206	
OLUMIANT]
ONGLYZA]
ONUREG		
OPSUMIT		
ORALAIR		
ORENCIA CLICKJECT		
ORENCIA IV		;
ORENCIA SC		;
ORENITRAM		;
ORILISSA		;
ORKAMBI		;
ORPHENADRINE		;
OSENI		;
OTEZLA		;
OXBRYTA		;
OXERVATE		;
PALYNZIQ		;
PEMAZYRE		;
PENNSAID		;
PERSERIS		;
PHENOBARBITAL		;
PIQRAY		;
PLEGRIDY		;
PRALUENT		;
PRETOMANID		;
PREVYMIS		;
PROCARDIA CAPSULES	233	1

PROLASTIN C	
PROMACTA	
РКОТОРІС	
PROVIGIL	
PURIXAN	
OINLOCK	
QUALAQUIN	
RADICAVA	
RAYALDEE	
REBLOZYL	
RELISTOR	
REMICADE	
RENFLEXIS	24
REPATHA	
RETEVMO	
REVATIO	
REVCOVI	25
REVLIMID	25
REXULTI	25
RINVOQ	25
RITUXIMAB	25
ROZLYTREK	25
RUBRACA	25
RUZURGI	25
RYDAPT	25
SAVELLA	26
SECUADO	26
SEROQUEL XR	26
SILIQ	26
SIMPONI(prefilled syringe, auto-injector)	26
SIMPONI ARIA	26
SKYRIZI	26
SOMA	
SOMAVERT	
SOVALDI	
SPRAVATO	
SPRITAM	
SPRYCEL	
STELARA IV	
STELARA SC	
STIVARGA	
STRENSIQ	
SUBSYS	
SUNOSI	27
SURMONTIL	~-

SYMLINPEN	.281
SYMPAZAN	282
TABRECTA	.283
TAFAMIDIS	284
TAGRISSO	285
TAKHZYRO	286
TALTZ	.287
TALZENNA	.288
TARCEVA	.289
TARGRETIN GEL	.290
TASIGNA	.291
TAVALISSE	292
TAZVERIK	293
TECENTRIQ	294
TECFIDERA	295
TEGSEDI	.296
TENEX	.297
TEPEZZA	.298
TETRABENAZINE	299
TIBSOVO	.300
TOLSURA	301
TREMFYA	.302
TRIHEXYPHENIDYL	.303
TRIKAFTA	304
TUKYSA	305
TURALIO	.306
TYMLOS	307
TYSABRI	.308
ULTRAVATE LOTION	.309
UPTRAVI	.310
VALCHLOR	311
VALTOCO	.312
VANCOCIN	.313
VENCLEXTA	314
VERSACLOZ	.315
VERZENIO	316
VIBERZI	317
VIMOVO	318
VINBLASTINE	319
VINCRISTINE	.320
VITRAKVI	.321
VIZIMPRO	.322
VOSEVI	.323
VOTRIENT	324
VRAYLAR	.325
VUMERITY	.326
VYONDYS 53	327

WAKIX	328
XALKORI	329
ХАТМЕР	
XCOPRI	331
XELJANZ	332
XENLETA	333
XEOMIN	334
XERMELO	
XOLAIR	
XOSPATA	337
XPOVIO	
XTANDI	339
YERVOY	
YONSA	341
ZALTRAP	342
ZARXIO	
ZEJULA	
ZELBORAF	345
ZEPATIER	
ZINPLAVA	347
ZULRESSO	348
ZYDELIG	
ZYKADIA	350
ZYTIGA	351